



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

Question 3.1:

What is the basic theme of organisation in the periodic table?

Answer

The basic theme of organisation of elements in the periodic table is to classify the elements in periods and groups according to their properties. This arrangement makes the study of elements and their compounds simple and systematic. In the periodic table, elements with similar properties are placed in the same group.

Question 3.2:

Which important property did Mendeleev use to classify the elements in his periodic table and did he stick to that?

Answer

Mendeleev arranged the elements in his periodic table ordered by atomic weight or mass. He arranged the elements in periods and groups in order of their increasing atomic weight. He placed the elements with similar properties in the same group.

However, he did not stick to this arrangement for long. He found out that if the elements were arranged strictly in order of their increasing atomic weights, then some elements did not fit within this scheme of classification.

Therefore, he ignored the order of atomic weights in some cases. For example, the atomic weight of iodine is lower than that of tellurium. Still Mendeleev placed tellurium (in Group **VI**) before iodine (in Group **VII**) simply because iodine's properties are so similar to fluorine, chlorine, and bromine.

Question 3.3:

What is the basic difference in approach between the Mendeleev's Periodic Law and the Modern Periodic Law?

Answer

Mendeleev's Periodic Law states that the physical and chemical properties of elements are periodic functions of their atomic weights. On the other hand, the Modern periodic Law states that the physical and chemical properties of elements are periodic functions of their atomic numbers.



Question 3.4:

On the basis of quantum numbers, justify that the sixth period of the periodic table should have 32 elements.

Answer

In the periodic table of the elements, a period indicates the value of the principal quantum number (n) for the outermost shells. Each period begins with the filling of principal quantum number (n). The value of n for the sixth period is 6. For $n = 6$, azimuthal quantum number (l) can have values of 0, 1, 2, 3, 4.

According to Aufbau's principle, electrons are added to different orbitals in order of their increasing energies. The energy of the $6d$ subshell is even higher than that of the $7s$ subshell.

In the 6th period, electrons can be filled in only $6s$, $4f$, $5d$, and $6p$ subshells. Now, $6s$ has one orbital, $4f$ has seven orbitals, $5d$ has five orbitals, and $6p$ has three orbitals. Therefore, there are a total of sixteen ($1 + 7 + 5 + 3 = 16$) orbitals available. According to Pauli's exclusion principle, each orbital can accommodate a maximum of 2 electrons. Thus, 16 orbitals can accommodate a maximum of 32 electrons.

Hence, the sixth period of the periodic table should have 32 elements.

Question 3.5:

In terms of period and group where would you locate the element with $Z = 114$?

Answer

Elements with atomic numbers from $Z = 87$ to $Z = 114$ are present in the 7th period of the periodic table. Thus, the element with $Z = 114$ is present in the 7th period of the periodic table.

In the 7th period, first two elements with $Z = 87$ and $Z = 88$ are s -block elements, the next 14 elements excluding $Z = 89$ i.e., those with $Z = 90 - 103$ are f -block elements, ten elements with $Z = 89$ and $Z = 104 - 112$ are d -block elements, and the elements with $Z = 113 - 118$ are p -block elements. Therefore, the element with $Z = 114$ is the second p -block element in the 7th period. Thus, the element with $Z = 114$ is present in the 7th period and 4th group of the periodic table.



Question 3.6:

Write the atomic number of the element present in the third period and seventeenth group of the periodic table.

Answer

There are two elements in the 1st period and eight elements in the 2nd period., The third period starts with the element with $Z = 11$. Now, there are eight elements in the third period. Thus, the 3rd period ends with the element with $Z = 18$ i.e., the element in the 18th group of the third period has $Z = 18$. Hence, the element in the 17th group of the third period has atomic number $Z = 17$.

Question 3.7:

Which element do you think would have been named by

- (i) Lawrence Berkeley Laboratory
- (ii) Seaborg's group?

Answer

- (i) Lawrencium (Lr) with $Z = 103$ and Berkelium (Bk) with $Z = 97$
- (ii) Seaborgium (Sg) with $Z = 106$

Question 3.8:

Why do elements in the same group have similar physical and chemical properties?

Answer

The physical and chemical properties of elements depend on the number of valence electrons. Elements present in the same group have the same number of valence electrons. Therefore, elements present in the same group have similar physical and chemical properties.

Question 3.9:

What does atomic radius and ionic radius really mean to you?

Answer



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

Atomic radius is the radius of an atom. It measures the size of an atom. If the element is a metal, then the atomic radius refers to the metallic radius, and if the element is a non-metal, then it refers to the covalent radius. Metallic radius is calculated as half the internuclear distance separating the metal cores in the metallic crystal. For example, the internuclear distance between two adjacent copper atoms in solid copper is 256 pm.

$$\frac{256}{2} \text{ pm} = 128 \text{ pm}$$

Thus, the metallic radius of copper is taken as

Covalent radius is measured as the distance between two atoms when they are found together by a single bond in a covalent molecule. For example, the distance between two chlorine atoms in chlorine molecule is 198 pm. Thus, the covalent radius of chlorine is

$$\frac{198}{2} \text{ pm} = 99 \text{ pm}$$

taken as

Ionic radius means the radius of an ion (cation or anion). The ionic radii can be calculated by measuring the distances between the cations and anions in ionic crystals.

Since a cation is formed by removing an electron from an atom, the cation has fewer electrons than the parent atom resulting in an increase in the effective nuclear charge.

Thus, a cation is smaller than the parent atom. For example, the ionic radius of Na^+ ion is 95 pm, whereas the atomic radius of Na atom is 186 pm. On the other hand, an anion is larger in size than its parent atom. This is because an anion has the same nuclear charge, but more electrons than the parent atom resulting in an increased repulsion among the electrons and a decrease in the effective nuclear charge. For example, the ionic radius of F^- ion is 136 pm, whereas the atomic radius of F atom is 64 pm.

Question 3.10:

How does atomic radius vary in a period and in a group? How do you explain the variation?

Answer

Atomic radius generally decreases from left to right across a period. This is because within a period, the outer electrons are present in the same valence shell and the atomic number increases from left to right across a period, resulting in an increased effective nuclear charge. As a result, the attraction of electrons to the nucleus increases.



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

On the other hand, the atomic radius generally increases down a group. This is because down a group, the principal quantum number (n) increases which results in an increase of the distance between the nucleus and valence electrons.

Question 3.11:

What do you understand by isoelectronic species? Name a species that will be isoelectronic with each of the following atoms or ions.

- (i) F^-
- (ii) Ar
- (iii) Mg^{2+}
- (iv) Rb^+

Answer

Atoms and ions having the same number of electrons are called isoelectronic species.

(i) F^- ion has $9 + 1 = 10$ electrons. Thus, the species isoelectronic with it will also have 10 electrons. Some of its isoelectronic species are Na^+ ion ($11 - 1 = 10$ electrons), Ne (10 electrons), O^{2-} ion ($8 + 2 = 10$ electrons), and Al^{3+} ion ($13 - 3 = 10$ electrons).

(ii) Ar has 18 electrons. Thus, the species isoelectronic with it will also have 18 electrons. Some of its isoelectronic species are S^{2-} ion ($16 + 2 = 18$ electrons), Cl^- ion ($17 + 1 = 18$ electrons), K^+ ion ($19 - 1 = 18$ electrons), and Ca^{2+} ion ($20 - 2 = 18$ electrons).

(iii) Mg^{2+} ion has $12 - 2 = 10$ electrons. Thus, the species isoelectronic with it will also have 10 electrons. Some of its isoelectronic species are F^- ion ($9 + 1 = 10$ electrons), Ne (10 electrons), O^{2-} ion ($8 + 2 = 10$ electrons), and Al^{3+} ion ($13 - 3 = 10$ electrons).

(iv) Rb^+ ion has $37 - 1 = 36$ electrons. Thus, the species isoelectronic with it will also have 36 electrons. Some of its isoelectronic species are Br^- ion ($35 + 1 = 36$ electrons), Kr (36 electrons), and Sr^{2+} ion ($38 - 2 = 36$ electrons).

Question 3.12:

Consider the following species:

N^{3-} , O^{2-} , F^- , Na^+ , Mg^{2+} and Al^{3+}



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

- (a) What is common in them?
(b) Arrange them in the order of increasing ionic radii.

Answer

(a) Each of the given species (ions) has the same number of electrons (10 electrons). Hence, the given species are isoelectronic.


(b) The ionic radii of isoelectronic species increases with a decrease in the magnitudes of nuclear charge.

The arrangement of the given species in order of their increasing nuclear charge is as follows:



Nuclear charge = +7 +8 +9 +11 +12 +13

Therefore, the arrangement of the given species in order of their increasing ionic radii is as follows:



Question 3.13:

Explain why cations are smaller and anions larger in radii than their parent atoms?

Answer

A cation has a fewer number of electrons than its parent atom, while its nuclear charge remains the same. As a result, the attraction of electrons to the nucleus is more in a cation than in its parent atom. Therefore, a cation is smaller in size than its parent atom. On the other hand, an anion has one or more electrons than its parent atom, resulting in an increased repulsion among the electrons and a decrease in the effective nuclear charge. As a result, the distance between the valence electrons and the nucleus is more in anions than in its parent atom. Hence, an anion is larger in radius than its parent atom.

Question 3.14:

What is the significance of the terms - 'isolated gaseous atom' and 'ground state' while defining the ionization enthalpy and electron gain enthalpy?

Hint: Requirements for comparison purposes.

Answer



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

Ionization enthalpy is the energy required to remove an electron from an isolated gaseous atom in its ground state. Although the atoms are widely separated in the gaseous state, there are some amounts of attractive forces among the atoms. To determine the ionization enthalpy, it is impossible to isolate a single atom. But, the force of attraction can be further reduced by lowering the pressure. For this reason, the term 'isolated gaseous atom' is used in the definition of ionization enthalpy.

Ground state of an atom refers to the most stable state of an atom. If an isolated gaseous atom is in its ground state, then less amount energy would be required to remove an electron from it. Therefore, for comparison purposes, ionization enthalpy and electron gain enthalpy must be determined for an 'isolated gaseous atom' and its 'ground state'.

Question 3.15:

Energy of an electron in the ground state of the hydrogen atom is -2.18×10^{-18} J. Calculate the ionization enthalpy of atomic hydrogen in terms of J mol^{-1} .

Answer

The energy of an electron in the ground state of the hydrogen atom is -2.18×10^{-18} J.

Therefore, the energy required to remove that electron from the ground state of hydrogen atom is 2.18×10^{-18} J.

$$\therefore \text{Ionization enthalpy of atomic hydrogen} = 2.18 \times 10^{-18} \text{ J}$$

Hence, ionization enthalpy of atomic hydrogen in terms of $\text{J mol}^{-1} = 2.18 \times 10^{-18} \times 6.02 \times 10^{23} \text{ J mol}^{-1} = 1.31 \times 10^6 \text{ J mol}^{-1}$

Question 3.16:

Among the second period elements the actual ionization enthalpies are in the order $\text{Li} < \text{B} < \text{Be} < \text{C} < \text{O} < \text{N} < \text{F} < \text{Ne}$.

Explain why

- (i) Be has higher $\Delta_i H$ than B
- (ii) O has lower $\Delta_i H$ than N and F?

Answer

(i) During the process of ionization, the electron to be removed from beryllium atom is a 2s-electron, whereas the electron to be removed from boron atom is a 2p-electron.



Now, $2s$ -electrons are more strongly attached to the nucleus than $2p$ -electrons. Therefore, more energy is required to remove a $2s$ -electron of beryllium than that required to remove a $2p$ -electron of boron. Hence, beryllium has higher $\Delta_i H$ than boron.

(ii) In nitrogen, the three $2p$ -electrons of nitrogen occupy three different atomic orbitals. However, in oxygen, two of the four $2p$ -electrons of oxygen occupy the same $2p$ -orbital. This results in increased electron-electron repulsion in oxygen atom. As a result, the energy required to remove the fourth $2p$ -electron from oxygen is less as compared to the energy required to remove one of the three $2p$ -electrons from nitrogen. Hence, oxygen has lower $\Delta_i H$ than nitrogen.

Fluorine contains one electron and one proton more than oxygen. As the electron is being added to the same shell, the increase in nuclear attraction (due to the addition of a proton) is more than the increase in electronic repulsion (due to the addition of an electron). Therefore, the valence electrons in fluorine atom experience a more effective nuclear charge than that experienced by the electrons present in oxygen. As a result, more energy is required to remove an electron from fluorine atom than that required to remove an electron from oxygen atom. Hence, oxygen has lower $\Delta_i H$ than fluorine.

Question 3.17:

How would you explain the fact that the first ionization enthalpy of sodium is lower than that of magnesium but its second ionization enthalpy is higher than that of magnesium?

Answer

The first ionization enthalpy of sodium is more than that of magnesium. This is primarily because of two reasons:

1. The atomic size of sodium is greater than that of magnesium
2. The effective nuclear charge of magnesium is higher than that of sodium

For these reasons, the energy required to remove an electron from magnesium is more than the energy required in sodium. Hence, the first ionization enthalpy of sodium is lower than that of magnesium.

However, the second ionization enthalpy of sodium is higher than that of magnesium. This is because after losing an electron, sodium attains the stable noble gas configuration. On the other hand, magnesium, after losing an electron still has one



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

electron in the 3s-orbital. In order to attain the stable noble gas configuration, it still has to lose one more electron. Thus, the energy required to remove the second electron in case of sodium is much higher than that required in case of magnesium. Hence, the second ionization enthalpy of sodium is higher than that of magnesium.

Question 3.18:

What are the various factors due to which the ionization enthalpy of the main group elements tends to decrease down a group?

Answer

The factors responsible for the ionization enthalpy of the main group elements to decrease down a group are listed below:

(i) Increase in the atomic size of elements: As we move down a group, the number of shells increases. As a result, the atomic size also increases gradually on moving down a group. As the distance of the valence electrons from the nucleus increases, the electrons are not held very strongly. Thus, they can be removed easily. Hence, on moving down a group, ionization energy decreases.

(ii) Increase in the shielding effect: The number of inner shells of electrons increases on moving down a group. Therefore, the shielding of the valence electrons from the nucleus by the inner core electrons increases down a group. As a result, the valence electrons are not held very tightly by the nucleus. Hence, the energy required to remove a valence electron decreases down a group.

Question 3.19:

The first ionization enthalpy values (in kJmol^{-1}) of group 13 elements are :

B	Al	Ga	In	Tl
801	577	579	558	589

How would you explain this deviation from the general trend?

Answer

On moving down a group, ionization enthalpy generally decreases due to an increase in the atomic size and shielding. Thus, on moving down group 13, ionization enthalpy decreases from B to Al. But, Ga has higher ionization enthalpy than Al. Al follows



immediately after s -block elements, whereas Ga follows after d -block elements. The shielding provided by d -electrons is not very effective. These electrons do not shield the valence electrons very effectively. As a result, the valence electrons of Ga experience a greater effective nuclear charge than those of Al. Further, moving from Ga to In, the ionization enthalpy decreases due to an increase in the atomic size and shielding. But, on moving from In to Tl, the ionization enthalpy again increases. In the periodic table, Tl follows after $4f$ and $5d$ electrons. The shielding provided by the electrons in both these orbitals is not very effective. Therefore, the valence electron is held quite strongly by the nucleus. Hence, the ionization energy of Tl is on the higher side.

Question 3.20:

Which of the following pairs of elements would have a more negative electron gain enthalpy?

(i) O or F (ii) F or Cl

Answer

(i) O and F are present in the same period of the periodic table. An F atom has one proton and one electron more than O and as an electron is being added to the same shell, the atomic size of F is smaller than that of O. As F contains one proton more than O, its nucleus can attract the incoming electron more strongly in comparison to the nucleus of O atom. Also, F needs only one more electron to attain the stable noble gas configuration. Hence, the electron gain enthalpy of F is more negative than that of O.

(ii) F and Cl belong to the same group of the periodic table. The electron gain enthalpy usually becomes less negative on moving down a group. However, in this case, the value of the electron gain enthalpy of Cl is more negative than that of F. This is because the atomic size of F is smaller than that of Cl. In F, the electron will be added to quantum level $n = 2$, but in Cl, the electron is added to quantum level $n = 3$. Therefore, there are less electron- electron repulsions in Cl and an additional electron can be accommodated easily. Hence, the electron gain enthalpy of Cl is more negative than that of F.

Question 3.21:

Would you expect the second electron gain enthalpy of O as positive, more negative or less negative than the first? Justify your answer.



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

Answer

When an electron is added to O atom to form O^- ion, energy is released. Thus, the first electron gain enthalpy of O is negative.



On the other hand, when an electron is added to O^- ion to form O^{2-} ion, energy has to be given out in order to overcome the strong electronic repulsions. Thus, the second electron gain enthalpy of O is positive.



Question 3.22:

What is the basic difference between the terms electron gain enthalpy and electronegativity?

Answer

Electron gain enthalpy is the measure of the tendency of an isolated gaseous atom to accept an electron, whereas electronegativity is the measure of the tendency of an atom in a chemical compound to attract a shared pair of electrons.

Question 3.23:

How would you react to the statement that the electronegativity of N on Pauling scale is 3.0 in all the nitrogen compounds?

Answer

Electronegativity of an element is a variable property. It is different in different compounds. Hence, the statement which says that the electronegativity of N on Pauling scale is 3.0 in all nitrogen compounds is incorrect. The electronegativity of N is different in NH_3 and NO_2 .

Question 3.24:

Describe the theory associated with the radius of an atom as it

- (a) gains an electron
- (b) loses an electron

Answer



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

(a) When an atom gains an electron, its size increases. When an electron is added, the number of electrons goes up by one. This results in an increase in repulsion among the electrons. However, the number of protons remains the same. As a result, the effective nuclear charge of the atom decreases and the radius of the atom increases.

(b) When an atom loses an electron, the number of electrons decreases by one while the nuclear charge remains the same. Therefore, the interelectronic repulsions in the atom decrease. As a result, the effective nuclear charge increases. Hence, the radius of the atom decreases.

Question 3.25:

Would you expect the first ionization enthalpies for two isotopes of the same element to be the same or different? Justify your answer.

Answer

The ionization enthalpy of an atom depends on the number of electrons and protons (nuclear charge) of that atom. Now, the isotopes of an element have the same number of protons and electrons. Therefore, the first ionization enthalpy for two isotopes of the same element should be the same.

Question 3.26:

What are the major differences between metals and non-metals?

Answer

	Metals		Non-metals
1.	Metals can lose electrons easily.	1.	Non-metals cannot lose electrons easily.
2.	Metals cannot gain electrons easily.	2.	Non-metals can gain electrons easily.
3.	Metals generally form ionic compounds.	3.	Non-metals generally form covalent compounds.
4.	Metals oxides are basic in nature.	4.	Non-metallic oxides are acidic in



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

			nature.
5.	Metals have low ionization enthalpies.	5.	Non-metals have high ionization enthalpies.
6.	Metals have less negative electron gain enthalpies.	6.	Non-metals have high negative electron gain enthalpies.
7.	Metals are less electronegative. They are rather electropositive elements.	7.	Non-metals are electronegative.
8.	Metals have a high reducing power.	8.	Non-metals have a low reducing power.

Question 3.27:

Use the periodic table to answer the following questions.

- Identify an element with five electrons in the outer subshell.
- Identify an element that would tend to lose two electrons.
- Identify an element that would tend to gain two electrons.
- Identify the group having metal, non-metal, liquid as well as gas at the room temperature.

Answer

(a) The electronic configuration of an element having 5 electrons in its outermost subshell should be $ns^2 np^5$. This is the electronic configuration of the halogen group. Thus, the element can be F, Cl, Br, I, or At.

(b) An element having two valence electrons will lose two electrons easily to attain the stable noble gas configuration. The general electronic configuration of such an element will be ns^2 . This is the electronic configuration of group 2 elements. The elements present in group 2 are Be, Mg, Ca, Sr, Ba.

(c) An element is likely to gain two electrons if it needs only two electrons to attain the stable noble gas configuration. Thus, the general electronic configuration of such an element should be $ns^2 np^4$. This is the electronic configuration of the oxygen family.

(d) Group 17 has metal, non-metal, liquid as well as gas at room temperature.

**Question 3.28:**

The increasing order of reactivity among group 1 elements is $\text{Li} < \text{Na} < \text{K} < \text{Rb} < \text{Cs}$ whereas that among group 17 elements is $\text{F} > \text{Cl} > \text{Br} > \text{I}$. Explain.

Answer

The elements present in group 1 have only 1 valence electron, which they tend to lose. Group 17 elements, on the other hand, need only one electron to attain the noble gas configuration. On moving down group 1, the ionization enthalpies decrease. This means that the energy required to lose the valence electron decreases. Thus, reactivity increases on moving down a group. Thus, the increasing order of reactivity among group 1 elements is as follows:



In group 17, as we move down the group from Cl to I, the electron gain enthalpy becomes less negative i.e., its tendency to gain electrons decreases down group 17. Thus, reactivity decreases down a group. The electron gain enthalpy of F is less negative than Cl. Still, it is the most reactive halogen. This is because of its low bond dissociation energy. Thus, the decreasing order of reactivity among group 17 elements is as follows:
 $\text{F} > \text{Cl} > \text{Br} > \text{I}$

Question 3.29:

Write the general outer electronic configuration of *s*-, *p*-, *d*- and *f*- block elements.

Answer

Element General outer electronic configuration

s-block ns^{1-2} , where $n = 2 - 7$

p-block ns^2np^{1-6} , where $n = 2 - 6$

d-block $(n-1)d^{1-10}ns^{0-2}$, where $n = 4 - 7$

f-block $(n-2)f^{1-14}(n-1)d^{0-10}ns^2$, where $n = 6 - 7$

Question 3.30:

Assign the position of the element having outer electronic configuration



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

(i) $ns^2 np^4$ for $n = 3$ (ii) $(n - 1)d^2 ns^2$ for $n = 4$, and (iii) $(n - 2)f^7 (n - 1)d^1 ns^2$ for $n = 6$, in the periodic table.

Answer

(i) Since $n = 3$, the element belongs to the 3rd period. It is a p -block element since the last electron occupies the p -orbital.

There are four electrons in the p -orbital. Thus, the corresponding group of the element
= Number of s -block groups + number of d -block groups + number of p -electrons
= 2 + 10 + 4
= 16

Therefore, the element belongs to the 3rd period and 16th group of the periodic table. Hence, the element is Sulphur.

(ii) Since $n = 4$, the element belongs to the 4th period. It is a d -block element as d -orbitals are incompletely filled.

There are 2 electrons in the d -orbital.

Thus, the corresponding group of the element
= Number of s -block groups + number of d -block groups
= 2 + 2
= 4

Therefore, it is a 4th period and 4th group element. Hence, the element is Titanium.

(iii) Since $n = 6$, the element is present in the 6th period. It is an f -block element as the last electron occupies the f -orbital. It belongs to group 3 of the periodic table since all f -block elements belong to group 3. Its electronic configuration is $[Xe] 4f^7 5d^1 6s^2$. Thus, its atomic number is $54 + 7 + 2 + 1 = 64$. Hence, the element is Gadolinium.

Question 3.31:

The first ($\Delta_i H_1$) and the second ($\Delta_i H$) ionization enthalpies (in kJ mol^{-1}) and the ($\Delta_{eg} H$) electron gain enthalpy (in kJ mol^{-1}) of a few elements are given below:

Elements	$\Delta_i H$	$\Delta_i H$	$\Delta_{eg} H$
I	520	7300	-60
II	419	3051	-48
III	1681	3374	-328
IV	1008	1846	-295



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

V	2372	5251	+48
VI	738	1451	-40

Which of the above elements is likely to be :

- (a) the least reactive element.
- (b) the most reactive metal.
- (c) the most reactive non-metal.
- (d) the least reactive non-metal.
- (e) the metal which can form a stable binary halide of the formula MX_2 , (X=halogen).
- (f) the metal which can form a predominantly stable covalent halide of the formula MX (X=halogen)?

Answer

- (a)** Element V is likely to be the least reactive element. This is because it has the highest first ionization enthalpy ($\Delta_i H_1$) and a positive electron gain enthalpy ($\Delta_{eg} H$).
- (b)** Element II is likely to be the most reactive metal as it has the lowest first ionization enthalpy ($\Delta_i H_1$) and a low negative electron gain enthalpy ($\Delta_{eg} H$).
- (c)** Element III is likely to be the most reactive non-metal as it has a high first ionization enthalpy ($\Delta_i H_1$) and the highest negative electron gain enthalpy ($\Delta_{eg} H$).
- (d)** Element V is likely to be the least reactive non-metal since it has a very high first ionization enthalpy ($\Delta_i H_2$) and a positive electron gain enthalpy ($\Delta_{eg} H$).
- (e)** Element VI has a low negative electron gain enthalpy ($\Delta_{eg} H$). Thus, it is a metal. Further, it has the lowest second ionization enthalpy ($\Delta_i H_2$). Hence, it can form a stable binary halide of the formula MX_2 (X=halogen).
- (f)** Element I has low first ionization energy and high second ionization energy. Therefore, it can form a predominantly stable covalent halide of the formula MX (X=halogen).

Question 3.32:

Predict the formula of the stable binary compounds that would be formed by the combination of the following pairs of elements.

- (a) Lithium and oxygen
- (b) Magnesium and nitrogen
- (c) Aluminium and iodine
- (d) Silicon and oxygen
- (e) Phosphorus and fluorine
- (f) Element 71 and fluorine



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

Answer

- (a) LiO_2
- (b) Mg_3N_2
- (c) AlI_3
- (d) SiO_2
- (e) PF_3 or PF_5
- (f) The element with the atomic number 71 is Lutetium (Lu). It has valency 3. Hence, the formula of the compound is LuF_3 .

Question 3.33:

In the modern periodic table, the period indicates the value of:

- (a) Atomic number
- (b) Atomic mass
- (c) Principal quantum number
- (d) Azimuthal quantum number.

Answer

The value of the principal quantum number (n) for the outermost shell or the valence shell indicates a period in the Modern periodic table.

Question 3.34:

Which of the following statements related to the modern periodic table is incorrect?

- (a) The p -block has 6 columns, because a maximum of 6 electrons can occupy all the orbitals in a p -shell.
- (b) The d -block has 8 columns, because a maximum of 8 electrons can occupy all the orbitals in a d -subshell.
- (c) Each block contains a number of columns equal to the number of electrons that can occupy that subshell.
- (d) The block indicates value of azimuthal quantum number (l) for the last subshell that received electrons in building up the electronic configuration.

Answer

The d -block has 10 columns because a maximum of 10 electrons can occupy all the orbitals in a d subshell.



Question 3.35:

Anything that influences the valence electrons will affect the chemistry of the element.

Which one of the following factors does not affect the valence shell?

- (a) Valence principal quantum number (n)
- (b) Nuclear charge (Z)
- (c) Nuclear mass
- (d) Number of core electrons.

Answer

Nuclear mass does not affect the valence electrons

Question 3.36:

The size of isoelectronic species — F^- , Ne and Na^+ is affected by

- (a) Nuclear charge (Z)
- (b) Valence principal quantum number (n)
- (c) Electron-electron interaction in the outer orbitals
- (d) None of the factors because their size is the same.

Answer

The size of an isoelectronic species increases with a decrease in the nuclear charge (Z).

For example, the order of the increasing nuclear charge of F^- , Ne, and Na^+ is as follows:



Z 9 10 11

Therefore, the order of the increasing size of F^- , Ne and Na^+ is as follows:



Question 3.37:

Which one of the following statements is incorrect in relation to ionization enthalpy?

- (a) Ionization enthalpy increases for each successive electron.
- (b) The greatest increase in ionization enthalpy is experienced on removal of electron from core noble gas configuration.
- (c) End of valence electrons is marked by a big jump in ionization enthalpy.



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

(d) Removal of electron from orbitals bearing lower n value is easier than from orbital having higher n value.

Answer

Electrons in orbitals bearing a lower n value are more attracted to the nucleus than electrons in orbitals bearing a higher n value. Hence, the removal of electrons from orbitals bearing a higher n value is easier than the removal of electrons from orbitals having a lower n value.

Question 3.38:

Considering the elements B, Al, Mg, and K, the correct order of their metallic character is:

- (a) $B > Al > Mg > K$ (b) $Al > Mg > B > K$
(c) $Mg > Al > K > B$ (d) $K > Mg > Al > B$

Answer

The metallic character of elements decreases from left to right across a period. Thus, the metallic character of Mg is more than that of Al.

The metallic character of elements increases down a group. Thus, the metallic character of Al is more than that of B.

Considering the above statements, we get $K > Mg$.

Hence, the correct order of metallic character is $K > Mg > Al > B$.

Question 3.39:

Considering the elements B, C, N, F, and Si, the correct order of their non-metallic character is:

- (a) $B > C > Si > N > F$ b) $Si > C > B > N > F$
(c) $F > N > C > B > Si$ d) $F > N > C > Si > B$

Answer

The non-metallic character of elements increases from left to right across a period. Thus, the decreasing order of non-metallic character is $F > N > C > B$.

Again, the non-metallic character of elements decreases down a group. Thus, the decreasing order of non-metallic characters of C and Si are $C > Si$. However, Si is less non-metallic than B i.e., $B > Si$.



Chapter 3 – Classification of Elements and Periodicity in Properties

Class XI

Chemistry

Hence, the correct order of their non-metallic characters is $F > N > C > B > Si$.

Question 3.40:

Considering the elements F, Cl, O and N, the correct order of their chemical reactivity in terms of oxidizing property is:

- (a) $F > Cl > O > N$ (b) $F > O > Cl > N$
(c) $Cl > F > O > N$ (d) $O > F > N > Cl$

Answer

The oxidizing character of elements increases from left to right across a period. Thus, we get the decreasing order of oxidizing property as $F > O > N$.

Again, the oxidizing character of elements decreases down a group. Thus, we get $F > Cl$. However, the oxidizing character of O is more than that of Cl i.e., $O > Cl$.

Hence, the correct order of chemical reactivity of F, Cl, O, and N in terms of their oxidizing property is $F > O > Cl > N$.

UNIT-3

CLASSIFICATION OF ELEMENTS AND PERIODICITY IN PROPERTIES

Moseley, the English physicist showed that atomic number is more fundamental property of an element than its atomic mass. Therefore, the position of an element in the periodic table depends on its atomic number.

Modern periodic laws : The physical and chemical properties of elements are the periodic functions of their atomic numbers.

Types of Elements : s-, p-, d- and f- blocks :

s-block elements : Group 1 (alkali metals) and group 2 elements (alkaline earth metals) which respectively have ns^1 and ns^2 outermost electronic configurations.

p-block elements belong to groups 13 to 18. The outermost electronic configuration is $ns^2 np^{1-6}$. He ($1s^2$) is a s-block element but is positioned with the group 18 elements ($ns^2 np^6$) because it has completely filled valence shell and as a result, exhibits properties characteristic of other noble gases.

d-block elements (Transition elements) are the elements of group 3 to 12 having outer electronic configuration $(n - 1)d^{1-10} ns^{1-2}$. Four transition series are 3d, 4d, 5d and 6d. 6d-series is incomplete. Atomic radius generally decreases across a period and increases as we descend the group.

A cation is smaller but the anion is larger than the parent atom. In case of isoelectronic species, the cation with greater positive charge has smaller radius but anion with greater negative charge has the larger radius.

Ionization enthalpy ($\Delta_i H$) is the enthalpy change for the reaction : $X_{(g)} \rightarrow X^+_{(g)} + e^-$

Second ionization enthalpy ($\Delta_i H_2$) is more than the first ionization enthalpy ($\Delta_i H_1$).

$$\Delta_i H_1 < \Delta_i H_2 < \Delta_i H_3 < \dots\dots\dots$$

For the elements of second period, the correct order of increasing ionization enthalpy ($\Delta_i H$) is :

$$\Delta_i H_1 : \text{Li} < \text{B} < \text{Be} < \text{C} < \text{O} < \text{N} < \text{F} < \text{Ne} ; \quad X(g) \rightarrow X(g)^+ + e^-$$

For the elements of third period, the correct increasing order of ionization enthalpy is :


$$\Delta_i H_1 : \text{Na} < \text{Al} < \text{Mg} < \text{Si} < \text{S} < \text{P} < \text{Cl} < \text{Ar}$$

For the group 1 elements, the ionization enthalpy decreases in the order :



Ionization enthalpy and atomic radius are closely related properties depending upon the attraction of electrons towards the nucleus and repulsion of electrons from each other.

Electron gain enthalpy ($\Delta_{\text{eg}}H$) is the enthalpy change for the reaction :



$\Delta_{\text{eg}}H$ for N atom is zero. All the elements except noble gases, have negative values of $\Delta_{\text{eg}}H$ whereas noble gases have positive values of $\Delta_{\text{eg}}H$.

***f*-block Elements (Inner-transition Series)**

Lanthanoides characterised by the filling of $4f$ orbitals, are the elements following lanthanum from $_{58}\text{Ce}$ to $_{71}\text{Lu}$. Actinoides characterised by the filling of $5f$ orbitals, are the elements following actinium from $_{70}\text{Th}$ to $_{103}\text{Lr}$. Characteristic outer electronic configuration is $(n-2)f^{1-14} (n-1)d^{0-1} ns^2$.

Cause of periodicity in the properties of elements is the periodic repetition of similar electronic configuration of elements as the atomic number increases. Regular variation in physical and chemical properties are observed as we descend the group and move across a period from left to right in the periodic table.

Trend in $\Delta_{\text{eg}}H$: Electron gain enthalpy generally becomes more negative across a period as we move right to left and within a group becomes less negative down the group.

Exceptions : P, S, Cl have more negative $\Delta_{\text{eg}}H$ than N, O, F respectively.

Second electron gain enthalpy of an atom is always positive.

Electronegativity (EN) : It is the qualitative measure of an atom in a chemical compound to attract the shared electrons to itself.

The most widely used scale is Pauling scale which is based on bond energy data. By giving a reference value of 2.1 to H, the maximum electronegativity value of 4 is assigned to F.

Screening effect – Decrease in the force of attraction exerted by the nucleus on the valency electrons due to the presence of electrons in the inner shells, is called screening effect or shielding effect.

Effective nuclear charge denoted by Z^* is given as

$$= \text{Nuclear Charge} - \text{Shielding effect}$$

The magnitude of effective nuclear charge increases in a period when we move from left to right but decreases down the group.

Anomalous Properties of Second Period Elements

Each element of second period, *i.e.*, first element of each of group 1 and 2 and groups 13-17 shows many properties which are not shown by its congeners. Their anomalous behaviour is attributed to their small size, large charge/radius ratio, high electronegativity, non-availability of *d*-orbitals in their valence shell. Thus the first member of each group has only four valence orbitals (one *2s* and three *2p* orbitals) for bonding, whereas the second member of the group has nine valence orbitals (one *3s*, three *3p* and five *3d* orbitals). As a consequence of this, maximum covalency of first member of each group is limited to '4', whereas the other members of the group can expand their valence shell to accommodate more than four pairs of electrons. For example, B from $[\text{BF}_4]^-$ and Al, $[\text{AlF}_6]^{3-}$. In addition to this, the first member of each group of *p*-block elements displays greater ability to form *pπ-pπ* multiple bonds to itself (e.g. $\text{C}=\text{C}$, $\text{C}\equiv\text{C}$, $\text{N}=\text{N}$, $\text{N}\equiv\text{N}$, $\text{O}=\text{O}$) and to other second period elements (e.g., $\text{C}=\text{O}$, $\text{C}=\text{N}$, $\text{C}\equiv\text{N}$, $\text{N}=\text{O}$) compared to subsequent members of the group.

1 - MARK QUESTIONS

1. Name the scientist who gave birth to the periodic table.
2. Name the property used by Mendeleev to classify the elements in his periodic table ?
3. State the Modern Periodic Law.
4. How many groups and periods are there in the long form of the periodic table ?
5. Write the IUPAC name and symbol for the element with atomic number 119.
6. How does ionization enthalpy vary in a group and in a period?
7. Arrange B, C, N, O elements in increasing order of electron gain enthalpy.
8. Write the electronic configuration of the element having atomic number 21.
9. Explain the term electron gain enthalpy.
10. Out of K and K^+ , which one would have larger size ?
11. Arrange the following elements in the increasing order of metallic character:
B, Al, Mg, K
12. Predict the position of the elements in the periodic table having the electronic configuration :

$$(n - 1)d^1 ns^2 \text{ for } n = 4$$

13. Among the elements of the second period Li to Ne, pick out the element :
- having the highest first ionization enthalpy.
 - having the highest electronegativity.
14. The first ($\Delta_i H_1$) and the second ($\Delta_i H_2$) ionization enthalpies (kJ mol^{-1}) of the three elements are given below :


	I	II	III
$\Delta_i H_1$	403	549	1142
$\Delta_i H_2$	2640	1060	2080

Identify the element which is likely to be :

- a non-metal.
- an alkaline earth metal.

[Ans. II – alkaline earth metal, III – non-metal]

15. Predict the formulae of the stable binary compounds that would be formed by the following pairs of elements :
- Aluminium and Bromine
 - Element ($Z = 120$) and Oxygen
16. The following species are isoelectronic with noble gas Argon :







Arrange them in order of increasing size.

[Hint : Greater the nuclear charge, smaller is the size.]

17. The ionic radius of F^- is 136 pm whereas the atomic radius of F is only 64 pm. Explain why ?
- [Hint : Size of an anion is larger than the parent atom]

18. Write the general electronic configuration of f -block elements.
19. Consider the elements N, P, O and S and arrange them in the order of increasing negative electron gain enthalpy.
20. Show by chemical reaction with water that K_2O is a basic oxide.
21. Elements A, B, C and D have the following electronic configurations :



Which among these will belong to the same group in the periodic table ?

22. Predict the position of element in the periodic table having the electronic configuration $(n-1) d^1 ns^2$ for $n = 4$.

2 - MARK QUESTIONS




- (a) State Newland's law of octaves.
(b) Write two anomalies of Mendeleev's periodic table.
- Name the groups of elements classified as *s*-, *p*-, *d*- and *f*-blocks in the modern periodic table.
- How are Li and Mg related to each other in the periodic table ? Write the name of another pair having such a relationship.
- (a) Name the first and last member of the *3d* series.
(b) To which block (*s*-, *p*-, *d*- or *f*-) does the element with atomic number 50 belong ?
- The ionization enthalpy per mole of atomic hydrogen is 1.313×10^6 J. Calculate the energy of the electron in the ground state of the hydrogen atom.

$$[\text{Hint : I.E./atom} = \frac{1.313 \times 10^6 \text{ J mol}^{-1}}{6.023 \times 10^{23} \text{ mol}^{-1}} = 2.18 \times 10^{-18} \text{ J}]$$

$$\text{I. E.} = E_{\infty} - E_1 \quad [\text{Ans. } E_1 = -2.18 \times 10^{-18} \text{ J}]$$

- An element belong to third period of *p*-block elements. It has four electrons in the outermost shell. Predict its group. How many unpaired electrons are present in an atom of that element ?
- Write the atomic number and electronic configuration of the elements of fourth period which has maximum number of unpaired electrons ?
- Why do the periodic properties such as ionization enthalpy, electron gain enthalpy and electronegativity shows a decreasing trend down the group but an increasing trend along a period ? Explain your answer.
- Identify the elements having the following description and write their electronic configuration also :
 - Group 14, period 3
 - Group 18, period 2
 - Group 1, period 6[Ans. (a) Si, (b) Ne, (c) Cs]
- On the basis of quantum numbers, justify that fifth period of the periodic table should have eighteen elements.
- Lanthanoids and actinoids are placed in separate rows at the bottom of the periodic table. Explain the reason for this arrangement.

12. The electronic configuration of three elements are given below :





- (a) Which element has the highest electron gain enthalpy ?
- (b) Which element has the lowest electron gain enthalpy ?
13. The element with atomic number 120 has not yet been discovered. In which group would you place this element ? Write its electronic configuration also.
14. Write the formulae of the oxides formed by sodium and chlorine and by means of their chemical reaction with water. Classify them as acidic or basic oxide.
15. Arrange the following species as stated, giving reasons :
- (a) Increasing order of electronegativity (H, F, Cl)
- (b) Increasing order of ionic size : N^{3-} , Na^+ , F^- , O^{2-} , Mg^{2+}
16. Write the oxidation state and covalency of Al in $[AlCl(H_2O)_5]^{2+}$.
17. Explain why cations are smaller than corresponding anions.
18. Why noble gases have largest atomic radii in their respective periods.
19. Which of the following will have the most negative electron gain enthalpy and which the least negative ?
- P, S, Cl, F
- Explain your answer.
20. Why is ionisation enthalpy of Boron
- (B) less than that of Beryllium (Be) and of oxygen.
- (O) is less than that of Nitrogen (N)?

3 - MARK QUESTIONS

1. (a) Name the scientist who discovered that atomic number is the more fundamental periodic property of an element than its atomic mass.
- (b) Mention three main features of the modern periodic table.
2. The diagram given below is a part of the periodic table. Study the table and answer the questions that follow :

- (c) The increasing order of reactivity among group 1 elements is $\text{Li} < \text{Na} < \text{K} < \text{Rb} < \text{Cs}$ whereas that among group 17 elements is $\text{F} > \text{Cl} > \text{Br} > \text{I}$.
8. Given below are the plots of first ionization enthalpy ($\Delta_i H$) of elements of the second period as a function of atomic number (Z) (Fig. 1) and $\Delta_i H$ of alkali metals as a function of Z (Fig. 2). Answer the questions that follows :



- (a) First ionization enthalpy generally increases as we go across a period and decreases as we descend a group.
- (b) Ionization enthalpy of boron is slightly less than that of beryllium.
- (c) Ionization enthalpy of nitrogen is more than that of oxygen.
9. A, B, C are three elements with atomic number $Z - 1$, Z and $Z + 1$ respectively. B is an inert gas. Answer the following questions :
- (a) Predict the groups of A and C.
- (b) Which out of the three elements has positive electron gain enthalpy ?
- (c) Which out of the three elements has the least value of ionization energy ?
10. The first ($\Delta_i H_1$) and the second ($\Delta_i H_2$) ionization enthalpies (in kJ/mol) and the ($\Delta_{eg} H$) electron gain enthalpy (in kJ/mol) of a few elements are given below :

Elements	$\Delta_i H_1$	$\Delta_i H_2$	$\Delta_{eg} H$
I	520	7300	-60
II	419	3051	-48
III	1681	3374	-328
IV	1008	1846	-295
V	2372	5251	+48
VI	738	1451	-40

Which of the above elements is likely to be :

- (a) the least reactive element.
- (b) the most reactive metal.
- (c) the most reactive non-metal.
- (d) the least reactive non-metal.
- (e) the metal which can form a stable binary halide of the formula MX_2 (X = halogen).
- (f) the metal which can form a predominately stable covalent halide of the formula MX (X = halogen).