

**Question 5.1:**

Write any two characteristics of Chemisorption.

Answer

1. Chemisorption is highly specific in nature. It occurs only if there is a possibility of chemical bonding between the adsorbent and the adsorbate.
2. Like physisorption, chemisorption also increases with an increase in the surface area of the adsorbent.

Question 5.2:

Why does physisorption decrease with the increase of temperature?

Answer

Physisorption is exothermic in nature. Therefore, in accordance with Le-Chatelieres's principle, it decreases with an increase in temperature. This means that physisorption occurs more readily at a lower temperature.

Question 5.3:

Why are powdered substances more effective adsorbents than their crystalline forms?

Answer

Powdered substances are more effective adsorbents than their crystalline forms because when a substance is powdered, its surface area increases and physisorption is directly proportional to the surface area of the adsorbent.

Question 5.4:

Why is it necessary to remove CO when ammonia is obtained by Haber's process?

Answer

It is important to remove CO in the synthesis of ammonia as CO adversely affects the activity of the iron catalyst, used in Haber's process.

Question 5.5:

Why is the ester hydrolysis slow in the beginning and becomes faster after sometime?

Answer

Ester hydrolysis can be represented as:



Ester + Water \longrightarrow Acid + Alcohol

The acid produced in the reaction acts as a catalyst and makes the reaction faster. Substances that act as catalysts in the same reaction in which they are obtained as products are known as autocatalysts.

Question 5.6:

What is the role of desorption in the process of catalysis?

Answer

The role of desorption in the process of catalysis is to make the surface of the solid catalyst free for the fresh adsorption of the reactants on the surface.

Question 5.7:

What modification can you suggest in the Hardy-Schulze law?

Answer

Hardy-Schulze law states that 'the greater the valence of the flocculating ion added, the greater is its power to cause precipitation.'

This law takes into consideration only the charge carried by an ion, not its size. The smaller the size of an ion, the more will be its polarising power. Thus, Hardy-Schulze law can be modified in terms of the polarising power of the flocculating ion. Thus, the modified Hardy-Schulze law can be stated as 'the greater the polarising power of the flocculating ion added, the greater is its power to cause precipitation.'

Question 5.8:

Why is it essential to wash the precipitate with water before estimating it quantitatively?

Answer

When a substance gets precipitated, some ions that combine to form the precipitate get adsorbed on the surface of the precipitate. Therefore, it becomes important to wash the precipitate before estimating it quantitatively in order to remove these adsorbed ions or other such impurities.



**Question 5.1:**

Distinguish between the meaning of the terms adsorption and absorption.

Give one example of each.

Answer

Adsorption is a surface phenomenon of accumulation of molecules of a substance at the surface rather than in the bulk of a solid or liquid. The substance that gets adsorbed is called the 'adsorbate' and the substance on whose surface the adsorption takes place is called the 'adsorbent'. Here, the concentration of the adsorbate on the surface of the adsorbent increases. In adsorption, the substance gets concentrated at the surface only. It does not penetrate through the surface to the bulk of the solid or liquid. For example, when we dip a chalk stick into an ink solution, only its surface becomes coloured. If we break the chalk stick, it will be found to be white from inside.

On the other hand, the process of *absorption* is a bulk phenomenon. In absorption, the substance gets uniformly distributed throughout the bulk of the solid or liquid.

Question 5.2:

What is the difference between physisorption and chemisorption?

Answer

	Physisorption	Chemisorption
1.	In this type of adsorption, the adsorbate is attached to the surface of the adsorbent with weak van der Waal's forces of attraction.	In this type of adsorption, strong chemical bonds are formed between the adsorbate and the surface of the adsorbent.
2.	No new compound is formed in the process.	New compounds are formed at the surface of the adsorbent.
3.	It is generally found to be reversible in nature.	It is usually irreversible in nature.



4.	Enthalpy of adsorption is low as weak van der Waal's forces of attraction are involved. The values lie in the range of 20-40 kJ mol ⁻¹ .	Enthalpy of adsorption is high as chemical bonds are formed. The values lie in the range of 40-400 kJ mol ⁻¹ .
5.	It is favoured by low temperature conditions.	It is favoured by high temperature conditions.
6.	It is an example of multi-layer adsorption	It is an example of mono-layer adsorption.

Question 5.3:

Give reason why a finely divided substance is more effective as an adsorbent.

Answer

Adsorption is a surface phenomenon. Therefore, adsorption is directly proportional to the surface area. A finely divided substance has a large surface area. Both physisorption and chemisorption increase with an increase in the surface area. Hence, a finely divided substance behaves as a good adsorbent.

Question 5.4:

What are the factors which influence the adsorption of a gas on a solid?

Answer

There are various factors that affect the rate of adsorption of a gas on a solid surface.

(1) Nature of the gas:

Easily liquefiable gases such as NH₃, HCl etc. are adsorbed to a great extent in comparison to gases such as H₂, O₂ etc. This is because Van der Waal's forces are stronger in easily liquefiable gases.

(2) Surface area of the solid

The greater the surface area of the adsorbent, the greater is the adsorption of a gas on the solid surface.

(3) Effect of pressure

Adsorption is a reversible process and is accompanied by a decrease in pressure. Therefore, adsorption increases with an increase in pressure.


**(4) Effect of temperature**

Adsorption is an exothermic process. Thus, in accordance with Le-Chatelier's principle, the magnitude of adsorption decreases with an increase in temperature.

Question 5.5:

What is an adsorption isotherm? Describe Freundlich adsorption isotherm.

Answer



The plot between the extent of adsorption $\left(\frac{x}{m}\right)$ against the pressure of gas (P) at constant temperature (T) is called the adsorption isotherm.

Freundlich adsorption isotherm:

Freundlich adsorption isotherm gives an empirical relationship between the quantity of gas adsorbed by the unit mass of solid adsorbent and pressure at a specific temperature.

From the given plot it is clear that at pressure P_s , $\frac{x}{m}$ reaches the maximum value. P_s is called the saturation pressure. Three cases arise from the graph now.

Case I- At low pressure:

The plot is straight and sloping, indicating that the pressure is directly proportional to

$$\frac{x}{m} \text{ i.e., } \frac{x}{m} \propto P.$$

$$\frac{x}{m} = kP \quad (k \text{ is a constant})$$

Case II- At high pressure:

When pressure exceeds the saturated pressure, $\frac{x}{m}$ becomes independent of P values.



$$\frac{x}{m} \propto P^0$$

$$\frac{x}{m} = k P^0$$

Case III- At intermediate pressure:

At intermediate pressure, $\frac{x}{m}$ depends on P raised to the powers between 0 and 1. This relationship is known as the Freundlich adsorption isotherm.

$$\frac{x}{m} \propto P^n$$


$$\frac{x}{m} = k P^n \quad n > 1$$

Now, taking log:

$$\log \frac{x}{m} = \log k + \frac{1}{n} \log P$$

On plotting the graph between $\log \left(\frac{x}{m} \right)$ and $\log P$, a straight line is obtained with the

slope equal to $\frac{1}{n}$ and the intercept equal to $\log k$.

**Question 5.6:**

What do you understand by activation of adsorbent? How is it achieved?

Answer

By activating an adsorbent, we tend to increase the adsorbing power of the adsorbent.

Some ways to activate an adsorbent are:



(i) By increasing the surface area of the adsorbent. This can be done by breaking it into smaller pieces or powdering it.

(ii) Some specific treatments can also lead to the activation of the adsorbent. For example, wood charcoal is activated by heating it between 650 K and 1330 K in vacuum or air. It expels all the gases absorbed or adsorbed and thus, creates a space for adsorption of gases.

Question 5.7:

What role does adsorption play in heterogeneous catalysis?

Answer

Heterogeneous catalysis:

A catalytic process in which the catalyst and the reactants are present in different phases is known as a heterogeneous catalysis. This heterogeneous catalytic action can be explained in terms of the adsorption theory. The mechanism of catalysis involves the following steps:

(i) Adsorption of reactant molecules on the catalyst surface.

(ii) Occurrence of a chemical reaction through the formation of an intermediate.

(iii) De-sorption of products from the catalyst surface

(iv) Diffusion of products away from the catalyst surface.

In this process, the reactants are usually present in the gaseous state and the catalyst is present in the solid state. Gaseous molecules are then adsorbed on the surface of the catalyst. As the concentration of reactants on the surface of the catalyst increases, the rate of reaction also increases. In such reactions, the products have very less affinity for the catalyst and are quickly desorbed, thereby making the surface free for other reactants.

Question 5.8:

Why is adsorption always exothermic?

Answer

Adsorption is always exothermic. This statement can be explained in two ways.

(i) Adsorption leads to a decrease in the residual forces on the surface of the adsorbent. This causes a decrease in the surface energy of the adsorbent. Therefore, adsorption is always exothermic.



(ii) ΔH of adsorption is always negative. When a gas is adsorbed on a solid surface, its movement is restricted leading to a decrease in the entropy of the gas i.e., ΔS is negative. Now for a process to be spontaneous, ΔG should be negative.

$$\therefore \Delta G = \Delta H - T\Delta S$$

Since ΔS is negative, ΔH has to be negative to make ΔG negative. Hence, adsorption is always exothermic.

Question 5.9:

How are the colloidal solutions classified on the basis of physical states of the dispersed phase and dispersion medium?

Answer

One criterion for classifying colloids is the physical state of the dispersed phase and dispersion medium. Depending upon the type of the dispersed phase and dispersion medium (solid, liquid, or gas), there can be eight types of colloidal systems.

	Dispersed phase	Dispersion medium	Type of colloid	Example
1.	Solid	Solid	Solid Sol	Gemstone
2.	Solid	Liquid	Sol	Paint
3.	Solid	Gas	Aerosol	Smoke
4.	Liquid	Solid	Gel	Cheese
5.	Liquid	Liquid	Emulsion	Milk
6.	Liquid	Gas	Aerosol	Fog
7.	Gas	Solid	Solid foam	Pumice stone
8.	Gas	Liquid	Foam	Froth

Question 5.10:

Discuss the effect of pressure and temperature on the adsorption of gases on solids.

Answer

**Effect of pressure**

Adsorption is a reversible process and is accompanied by a decrease in pressure. Therefore, adsorption increases with an increase in pressure.

Effect of temperature

Adsorption is an exothermic process. Thus, in accordance with Le-Chatelier's principle, the magnitude of adsorption decreases with an increase in temperature.

Question 5.11:

What are lyophilic and lyophobic sols? Give one example of each type. Why are hydrophobic sols easily coagulated?

Answer

(i) Lyophilic sols:

Colloidal sols that are formed by mixing substances such as gum, gelatin, starch, etc. with a suitable liquid (dispersion medium) are called lyophilic sols. These sols are reversible in nature i.e., if two constituents of the sol are separated by any means (such as evaporation), then the sol can be prepared again by simply mixing the dispersion medium with the dispersion phase and shaking the mixture.

(ii) Lyophobic sols:

When substances such as metals and their sulphides etc. are mixed with the dispersion medium, they do not form colloidal sols. Their colloidal sols can be prepared only by special methods. Such sols are called lyophobic sols. These sols are irreversible in nature. For example: sols of metals.

Now, the stability of hydrophilic sols depends on two things- the presence of a charge and the salvation of colloidal particles. On the other hand, the stability of hydrophobic sols is only because of the presence of a charge. Therefore, the latter are much less stable than the former. If the charge of hydrophobic sols is removed (by addition of electrolytes), then the particles present in them come closer and form aggregates, leading to precipitation.

Question 5.12:

What is the difference between multimolecular and macromolecular colloids? Give one example of each. How are associated colloids different from these two types of colloids?

Answer



(i) In multi-molecular colloids, the colloidal particles are an aggregate of atoms or small molecules with a diameter of less than 1 nm. The molecules in the aggregate are held together by van der Waal's forces of attraction. Examples of such colloids include gold sol and sulphur sol.

(ii) In macro-molecular colloids, the colloidal particles are large molecules having colloidal dimensions. These particles have a high molecular mass. When these particles are dissolved in a liquid, sol is obtained. For example: starch, nylon, cellulose, etc.


(iii) Certain substances tend to behave like normal electrolytes at lower concentrations. However, at higher concentrations, these substances behave as colloidal solutions due to the formation of aggregated particles. Such colloids are called aggregated colloids.

Question 5.13:

What are enzymes? Write in brief the mechanism of enzyme catalysis.

Answer


Enzymes are basically protein molecules of high molecular masses. These form colloidal solutions when dissolved in water. These are complex, nitrogenous organic compounds produced by living plants and animals. Enzymes are also called 'biochemical catalysts'.



Mechanism of enzyme catalysis:

On the surface of the enzymes, various cavities are present with characteristic shapes. These cavities possess active groups such as $-\text{NH}_2$, $-\text{COOH}$, etc. The reactant molecules having a complementary shape fit into the cavities just like a key fits into a lock. This leads to the formation of an activated complex. This complex then decomposes to give the product.

Hence,



(Activated complex)



Step 2: $ES^+ \rightarrow E + P$

Question 5.14:

How are colloids classified on the basis of

- (i) Physical states of components
- (ii) Nature of dispersion medium and
- (iii) Interaction between dispersed phase and dispersion medium?

Answer

Colloids can be classified on various bases:

(i) On the basis of the physical state of the components (by components we mean the dispersed phase and dispersion medium). Depending on whether the components are solids, liquids, or gases, we can have eight types of colloids.

(ii) On the basis of the dispersion medium, sols can be divided as:

Dispersion medium	Name of sol
Water	Aquasol or hydrosol
Alcohol	Alcosol
Benzene	Benzosol
Gases	Aerosol

(iii) On the basis of the nature of the interaction between the dispersed phase and dispersion medium, the colloids can be classified as lyophilic (solvent attracting) and lyophobic (solvent repelling).

Question 5.15:

Explain what is observed

- (i) When a beam of light is passed through a colloidal sol.
- (ii) An electrolyte, NaCl is added to hydrated ferric oxide sol.
- (iii) Electric current is passed through a colloidal sol?

Answer



(i) When a beam of light is passed through a colloidal solution, then scattering of light is observed. This is known as the Tyndall effect. This scattering of light illuminates the path of the beam in the colloidal solution.

(ii) When NaCl is added to ferric oxide sol, it dissociates to give Na^+ and Cl^- ions. Particles of ferric oxide sol are positively charged. Thus, they get coagulated in the presence of negatively charged Cl^- ions.

(iii) The colloidal particles are charged and carry either a positive or negative charge. The dispersion medium carries an equal and opposite charge. This makes the whole system neutral. Under the influence of an electric current, the colloidal particles move towards the oppositely charged electrode. When they come in contact with the electrode, they lose their charge and coagulate.

Question 5.16:

What are emulsions? What are their different types? Give example of each type.

Answer

The colloidal solution in which both the dispersed phase and dispersion medium are liquids is called an emulsion.

There are two types of emulsions:

(a) Oil in water type:

Here, oil is the dispersed phase while water is the dispersion medium. For example: milk, vanishing cream, etc.

(b) Water in oil type:

Here, water is the dispersed phase while oil is the dispersion medium. For example: cold cream, butter, etc.

Question 5.17:

What is demulsification? Name two demulsifiers.

Answer

The process of decomposition of an emulsion into its constituent liquids is called demulsification. Examples of demulsifiers are surfactants, ethylene oxide, etc.

Question 5.18:

Action of soap is due to emulsification and micelle formation. Comment.



Answer

The cleansing action of soap is due to emulsification and micelle formation. Soaps are basically sodium and potassium salts of long chain fatty acids, $R\text{-COO}^-\text{Na}^+$. The end of the molecule to which the sodium is attached is polar in nature, while the alkyl-end is non-polar. Thus, a soap molecule contains a hydrophilic (polar) and a hydrophobic (non-polar) part.


When soap is added to water containing dirt, the soap molecules surround the dirt particles in such a manner that their hydrophobic parts get attached to the dirt molecule and the hydrophilic parts point away from the dirt molecule. This is known as micelle formation. Thus, we can say that the polar group dissolves in water while the non-polar group dissolves in the dirt particle. Now, as these micelles are negatively charged, they do not coalesce and a stable emulsion is formed.

Question 5.19:


Give four examples of heterogeneous catalysis.

Answer

(i) Oxidation of sulphur dioxide to form sulphur trioxide. In this reaction, Pt acts as a catalyst.




(ii) Formation of ammonia by the combination of dinitrogen and dihydrogen in the presence of finely divided iron.




This process is called the Haber's process.

(iii) Oswald's process: Oxidation of ammonia to nitric oxide in the presence of platinum.



(iv) Hydrogenation of vegetable oils in the presence of Ni.



Question 5.20:

What do you mean by activity and selectivity of catalysts?






Answer

(a) Activity of a catalyst:

The activity of a catalyst is its ability to increase the rate of a particular reaction. Chemisorption is the main factor in deciding the activity of a catalyst. The adsorption of reactants on the catalyst surface should be neither too strong nor too weak. It should just be strong enough to make the catalyst active.

(b) Selectivity of the catalyst:

The ability of the catalyst to direct a reaction to yield a particular product is referred to as the selectivity of the catalyst. For example, by using different catalysts, we can get different products for the reaction between H_2 and CO .



Question 5.21:

Describe some features of catalysis by zeolites.

Answer

Zeolites are alumino-silicates that are micro-porous in nature. Zeolites have a honeycomb-like structure, which makes them shape-selective catalysts. They have an extended 3D-network of silicates in which some silicon atoms are replaced by aluminium atoms, giving them an Al–O–Si framework. The reactions taking place in zeolites are very sensitive to the pores and cavity size of the zeolites. Zeolites are commonly used in the petrochemical industry.

Question 5.22:

What is shape selective catalysis?

Answer

A catalytic reaction which depends upon the pore structure of the catalyst and on the size of the reactant and the product molecules is called shape-selective catalysis. For example, catalysis by zeolites is a shape-selective catalysis. The pore size present in the



zeolites ranges from 260-740 pm. Thus, molecules having a pore size more than this cannot enter the zeolite and undergo the reaction.

Question 5.23:

Explain the following terms:

- (i)** Electrophoresis **(ii)** Coagulation
(iii) Dialysis **(iv)** Tyndall effect.

Answer

(i) Electrophoresis:

The movement of colloidal particles under the influence of an applied electric field is known as electrophoresis. Positively charged particles move to the cathode, while negatively charged particles move towards the anode. As the particles reach oppositely charged electrodes, they become neutral and get coagulated.

(ii) Coagulation:

The process of settling down of colloidal particles i.e., conversion of a colloid into a precipitate is called coagulation.

(iii) Dialysis

The process of removing a dissolved substance from a colloidal solution by the means of diffusion through a membrane is known as dialysis. This process is based on the principle that ions and small molecules can pass through animal membranes unlike colloidal particles.

(iv) Tyndall effect:

When a beam of light is allowed to pass through a colloidal solution, it becomes visible like a column of light. This is known as the Tyndall effect. This phenomenon takes place as particles of colloidal dimensions scatter light in all directions.

Question 5.24:

Give four uses of emulsions.

Answer

Four uses of emulsions:

- (i)** Cleansing action of soaps is based on the formation of emulsions.
(ii) Digestion of fats in intestines takes place by the process of emulsification.
(iii) Antiseptics and disinfectants when added to water form emulsions.




(iv) The process of emulsification is used to make medicines.

Question 5.25:

What are micelles? Give an example of a micellers system.

Answer

Micelle formation is done by substances such as soaps and detergents when dissolved in water. The molecules of such substances contain a hydrophobic and a hydrophilic part. When present in water, these substances arrange themselves in spherical structures in such a manner that their hydrophobic parts are present towards the centre, while the hydrophilic parts are pointing towards the outside (as shown in the given figure). This is known as micelle formation.



Question 5.26:

Explain the terms with suitable examples:

(i) Alcosol (ii) Aerosol (iii) Hydrosol

Answer

(i) Alcosol:

A colloidal solution having alcohol as the dispersion medium and a solid substance as the dispersed phase is called an alcosol.

For example: colloidal sol of cellulose nitrate in ethyl alcohol is an alcosol.

(ii) Aerosol:



A colloidal solution having a gas as the dispersion medium and a solid as the dispersed phase is called an aerosol.

For example: fog

(iii) Hydrosol

A colloidal solution having water as the dispersion medium and a solid as the dispersed phase is called a hydrosol.

For example: starch sol or gold sol

Question 5.27:

Comment on the statement that “colloid is not a substance but a state of substance”.

Answer

Common salt (a typical crystalloid in an aqueous medium) behaves as a colloid in a benzene medium. Hence, we can say that a colloidal substance does not represent a separate class of substances. When the size of the solute particle lies between 1 nm and 1000 nm, it behaves as a colloid.

Hence, we can say that colloid is not a substance but a state of the substance which is dependent on the size of the particle. A colloidal state is intermediate between a true solution and a suspension.

Experiment	Time in S	Total Pressure in atm
1	0	0.4
2	100	0.7

Calculate the rate constant. (Given : $\log 4 = 0.6021$, $\log 2 = 0.3010$)

Ans. $k = \frac{2.303}{t} \log \frac{P_i}{2P_i - P_t}$

$$k = \frac{2.303}{100} \log \frac{0.4}{2 \times 0.4 - 0.7}$$

$$= \frac{2.303 \times 0.6021}{100} = 0.01387 \text{ s}^{-1}$$

CHAPTER-5 Surface Chemistry

4 Marks Chapter

The branch of the Chemistry which deals with the study of surface phenomena is called surface Chemistry.

1. Adsorption: - The accumulation of molecules species at the surface rather in the bulk of a solid or liquid is termed adsorption.

2. Desorption:-Removal of adsorbate from the surface of adsorbent is known as Desorption.

3. Sorption:-When adsorption and absorption both takes place simultaneously.

4. Type of adsorption: - On the basis of interaction between adsorption and absorption, adsorbate are of two types:

(i)**Physical adsorption**/physisorption: - When weak Vander Waal's interaction involve between adsorbate and adsorbent.

(ii) **Chemical adsorption**/chemisorption:-When chemical bonds form between adsorbate and adsorbent.

5. Adsorption isotherm:-The variation in the amount of gas adsorbed by the adsorbent with pressure at constant temperature can be expressed by means of a curve termed as adsorption isotherm.


6. Application of adsorption:-

(a) Removal of colouring matter from solution using animal charcoal.

(b) Creating high vacuum

(c) In coal mines activated charcoal is used in gas mask to remove poisonous gases

7. Freundlich adsorption isotherm:-It is a graph which shows relationship between the quantity of gas adsorbed by unit mass of solid adsorbent and pressure at a particular temperature.



Pressure

$$x/m = kp^{1/n} \text{ ----- (1)}$$

$$\log (x/m) = \log K + 1/n \log P \text{ ----- (2)}$$

8 .Factors affecting adsorption:-

(i)Surface area: - Adsorption increases with increases of surface area of adsorbent.

(ii)Nature of adsorbate:- Easily liquefiable gases are readily adsorbed.


(iii)Temperature:-Low temperature is favorable for physical adsorption and High temperature for chemisorption.

(iv)Pressure: - Pressure increases, adsorption increases.


9. **CATALYSIS:-** Substances which alter the rate of chemical reaction and themselves remain chemically and quantitatively unchanged after the reaction are known as catalyst and the phenomenon is known as catalysis.

10. PROMOTERS AND POISONS

Promoters are substance that enhance the activity of catalyst while poisons decrease the activity of catalyst.



11. **Homogenous catalyst** – when reactants and catalyst are in same phase.



12. **Heterogeneous catalyst** - the catalytic process in which the reactants and catalyst are in different phase.




13. **Adsorption theory of Heterogeneous catalysis** – It explains the mechanism of heterogeneous catalyst.

The mechanism involves 5 steps:-


- a. Diffusion of reactants to the surface of catalyst.
- b. Adsorption of reactant molecules on the surface of catalyst.
- c. Chemical reaction on the catalyst surface through formation of an intermediate.
- d. Desorption of reaction product from the catalyst surface.
- e. Diffusion of reaction product away from the catalyst surface.

14. **IMPORTANT FEATURES OF SOLID CATALYST**

Activity - The activity of a catalyst depend on the strength of chemisorption. Catalytic activity increases from group 5 to group 11 elements of the periodic table.



Selectivity – The selectivity of a catalyst is its ability to direct a reaction to yield a particular product.



15. **SHAPE SELECTIVE CATALYSIS**

The catalytic reaction that depends upon the pure substance of the catalyst and the size of reactant and product molecules is called shape selective catalysis.

e.g. Zeolites are good shape selective catalyst.

ZSM-5 converts alcohol to gasoline.

16. **ENZYME CATALYSIS**

Enzymes are protein molecules of high molecular mass which catalyse the biochemical reaction.

e.g. Inversion of cane sugar by invertase enzyme.

17. **Characteristic of enzyme catalysis** –

- a. Enzymes are specific to substrate.
- b. Enzymes are highly active under optimum temperature.
- c. Enzymes are specific to pH. e.g. Pepsin act in acidic medium
- d. Enzymes are inhibited by the presence of certain substance.

Mechanism of enzyme catalysis –

1. Binding of enzyme to substrate to form an activated complex.



2. Decomposition of activated complex to form product.



18. **Colloid**- a colloid is a heterogeneous system in which one substance is dispersed (dispersed phase) in another substance called dispersion medium and size of dispersed phase is from 1nm-1000 nm.

19. TYPES OF COLLOIDS

(1) On the basis of nature of interaction between dispersed phase and dispersion medium.

(a) Lyophobic colloid-solvent hating colloid, these colloids cannot be prepared by simply mixing of dispersed phase into dispersion medium.

e.g. metallic sols.

(b) Lyophobic colloid-solvent loving these colloids can be prepared by simply mixing of dispersion phase into dispersion medium.

e.g. Starch sol.

(2) On the basis of types of particles of the dispersed phase

(a) Multimolecular colloid-on dissolution, a large number of atoms or smaller molecules of a substance aggregate together to form species having size in colloidal range. The species thus formed are called Multimolecular colloids.

e.g. Sulphur sol.

(b) Macromolecular colloids -macromolecules are suitable solvent from solution in which size of the particles are in range of colloidal range.

e.g. starch sol.

(c) Associated colloids (micelles)-some substances in low concentration behaves as normal strong electrolyte but at higher concentration exhibit colloidal behavior due to formation of aggregates. The aggregated particles are called micelles and also known as associated colloids.

(3) Kraft temperature- Temp. above which formation of micelles takes places.

(4) Critical micelle concentration (cmc) - concentration above which micelle formation takes place is known as cmc.


(5) PREPERATION OF COLLOIDS

(a) Chemical methods- By double decomposition, oxidation reaction or hydrolysis

OXIDATION



HYDROLYSIS



(sol)

(b) Bredig's arc method- For preparation of metallic sol. It involves dispersion as well as condensation.

(c) Peptization- Process of converting a precipitate into colloidal sol. By shaking it with dispersion medium in the presence of a small amount of electrolyte.

(6) PURIFICATION OF COLLOIDAL SOLUTION :-

(a) Dialysis-it is a process of removing a dissolved substance from a colloidal solution by membrane.

(b) Electro dialysis-when dialysis is carried out with an electric field applied around the membrane.

(c) Ultra filtration- Use of special filters which are permeable to all ionic substances except colloidal particles.

(7) PROPERTIES OF COLLOIDAL SOLUTION:-

(1) They show colligative properties

(2) Brownian movement-zig-zag motion of colloidal particles

(3) Tyndall effect-scattering of light by colloidal particles by which path of beam becomes clearly visible. This effect is known as tyndall effect.

1. Charge on colloidal particles – Colloidal particles which carry on electric charge and nature of charge is same on all particles.

2. Electrophoresis - Movement of Colloidal particles towards opposite electrode in presence of external electric field.

3. Coagulation – The process of setting of colloidal particles is called coagulation of the sol.

4. Hardy Schultz's Law – Coagulating power of a coagulating ion is directly proportional to the charge on the ion.

Eg: $\text{Na}^+ < \text{Ca}^{++} < \text{Al}^{3+}$ for negatively charged sol.

$\text{Cl}^- < \text{CO}_3^{2-} < \text{PO}_4^{3-} < [\text{Fe (CN)}_6]^{4-}$ for positive sol.

5. Emulsion – Liquid – liquid colloidal system is known as Emulsion.

There are two types of Emulsion.

- a) O/W type - Oil dispersed in water. Eg: milk, vanishing cream.
 b) W/O type - Water dispersed in oil. Eg: Butter & Cream.
6. Emulsifying Agent - The substance which stabilizes emulsion.

VERY SHORT ANSWER TYPE QUESTION

(1 marks)

- 1.** What are the physical states of dispersed phase and dispersion medium of froth?

Ans - Dispersed phase is gas, dispersion medium is liquid.

- 2.** What is the cause of Brownian movement among colloidal particles?

Ans - Charge on the colloidal particles due to which they suffer elastic collisions.

- 3.** Arrange the solutions: True solution, colloidal solution, suspension in decreasing order of their particles size?

Ans - Suspension > colloidal > true solution.

- 4.** Why is it necessary to remove CO when ammonia is obtained by Haber's process?

Ans- CO acts as poison catalyst for Haber's process therefore it will lower the activity of solution therefore it is necessary to remove when NH₃ obtained by Haber's process.

- 5.** How is adsorption of a gases related to its critical temperature?

Ans- Higher the critical temperature of the gas. Greater is the ease of liquefaction.

i.e. greater Vander walls forces of attraction and hence large adsorption will occur.

- 6.** What is meant by Shape Selective Catalyst?

Ans - On the Shape Selective Catalyst, the rate depends upon pore size of the catalyst and the shape & size of the reactant and products molecules.

- 7.** Of the physisorption & chemisorptions, which type of adsorption has higher enthalpy of adsorption?and why?

Ans - chemisorptions.Because of the chemical bonding between adsorbate and adsorbent.

- 8.** Give one example each of oil in water and water in oil emulsion. **CBSE-2014**

Ans: Oil in water : Milk , Water in oil: Butter

SHORT ANSWER TYPE QUESTION

(2 marks)

- 1.** Differentiate between physical & chemical adsorption?

Ans -

Physical adsorption	Chemical adsorption
a) Forces between adsorbate & adsorbent are weak Vander waal forces.	a) Forces between adsorbate & adsorbent are strong chemical forces.
b) Low heat of Adsorption.	b) High heat of Adsorption.

- 2.** Differentiate between Lyophobic & Lyophilic colloids?

3. Ans -

Lyophilic colloids	Lyophobic colloids
a) These are easily formed by direct mixing.	a) These are easily formed by Special method.

b) Particles of colloids are not easily visible even under ultra microscope. c) These are very stable.	b) Particles of colloids are easily visible under ultra microscope. c) These are unstable.
---	---

4. Differentiate between multi molecular, macromolecular and associated colloids?

5. Ans:-

Multi molecular colloids	Macromolecular colloids	Associated colloids
a) They consist of aggregates of atoms or molecules which generally have diameter less than 1nm. b) They are usually lyophobic	a) They consist of large molecules. b) They are hydrophilic.	a) Behave as colloidal size particles at higher conc. b) They have both lyophobic character & Lyophilic

6. What is difference between Sol. & Gel?

Ans – Both are colloidal solutions. Sol has solid as ‘dispersed phase & liquid as dispersion medium’. While in ‘Gel’ liquid is dispersed phase and solid is dispersion medium.


SHORT ANSWER TYPE QUESTION

(3 marks)

1. Discuss the effect of pressure & temperature on the adsorption of gases on solids?

Ans – Effect of pressure on adsorption: - At constant temp the extent of adsorption of gas(x/m) in the solid increases with pressure. A graph between x/m and the pressure P of a gas at constant temp is called adsorption isotherm.

2. See the diagram and answer:



Excess AgNO₃

Excess KI

- (1) Which type of colloid will form in both the cases? **(+ve and – ve respectively)**
(2) What is the cause of the charge on colloidal particles? **(adsorption of ions on ppt)**
(3) Name the process by which the colloid in this way form. **(Peptization)**

3. Explain What is observed when

- i) An electrolyte, NaCl is added to hydrate ferric oxide sol.
ii) Electric current is passed through a colloidal sol.
iii) When a beam of light is passed through a colloidal sol.

Ans –(i) The positively charged colloidal particles of Fe(OH)₃ get coagulated by the positively charged Cl⁻ ions provided by NaCl.

(ii) On passing direct current, colloidal particles move towards the positively charged electrode where they lose their charge and get coagulated.

(iii) Scattering of light by the colloidal particles takes place and the path of light becomes visible (Tyndall effect).

4. Describes some features of catalysis by Zeolites?

Ans – Features of catalysis by Zeolites:-

- I)** Zeolites are hydrated alumino silicates which have a three dimensional network structure containing water molecules in their pores.
II) To use them as catalysts, they heated so that water of hydration present in the pores is lost and the pores become vacant.
III) The size of pores varies from 260 to 740 pm. Thus, only those molecules can be adsorbed in these pores and catalyzed whose size is small enough to enter these pores. Hence, they act as molecular sieves or shape selective catalysts.

An important catalyst used in petroleum industries in zsm-5. It converts alcohols into petrol by first dehydrating them to form a mixture of hydro carbons.

Alcohols $\xrightarrow[\text{Dehydration}]{\text{ZSm-5}}$ Hydro carbons

5. Comment on the statement that “colloid is not a substance but state of a substance”?

Ans – The given statement is true. This is because the statement may exist as a colloid under certain conditions and as a crystalloid under certain other conditions. e.g. NaCl in water behaves as a crystalloid while in benzene, behaves as a colloid (called associated colloid). It is the size of the particles which matters i.e. the state in which the substance exist. If the size of the particles lies in the range 1nm to 1000nm it is in the colloid state.

6. Write short notes on followings:-

- a. Tyndall effect
- b. Brownian Movement
- c. Hardy Schulze Rule

Ans- (a) Tyndall effect-scattering of light by colloidal particles by which path of beam becomes clearly visible. this effect is known as tyndall effect

(b) Brownian movement-zig-zag motion of colloidal particles.

(c) Hardy Schulze Law – Coagulating value of a coagulating ion is directly proportional to the charge on the ion.

e.g: $\text{Na}^+ < \text{Ca}^{+2} < \text{Al}^{3+}$ for negatively charged sol.

$\text{Cl}^{-1} < \text{CO}_3^{2-} < \text{PO}_4^{3-} < [\text{Fe}(\text{CN})_6]^{4-}$ for positive sol.

7. a. In reference to Freundlich adsorption isotherm, write the expression for adsorption of gases on solids in the form of an equation.

CBSE-2014

b. Write an important characteristic of lyophilic sols.

c. Based on the type of particles dispersed phase, give one example each of associated colloid and multimolecular colloid.

Ans: a) $x/m = kP^{1/n}$

b. Lyophilic sols are reversible sols. These are quite stable and can not be coagulated.

c. Associated colloid : Soap solution, detergent
Multimolecular colloid: Sulphur sol, gold sol

8. Give reason:

- (a) Delta is formed where sea and river water meet.
- (b) To purify water alum is mixed in it.
- (c) Alum is applied to stop bleeding.

8. What happens when?

- (a) A colloid is repeatedly dialyzed.
- (b) A beam of light is passed through milk.
- (c) Electricity is passed through a colloid.

9. Explain following:

- (a) Peptization
- (b) Coagulation
- (c) Zeolites

10. What is the use of following?

- (1) Activated Charcoal
- (2) Silica gel
- (3) Coconut charcoal

11. Write three applications of adsorption.

12. Write three applications of colloid.

Chapter:-6 General Principles & Process of Isolation of Elements

3 Marks Chapter

Important Points :

1. The chemical substances in the earth's crust obtained by mining are called Minerals.
2. Minerals, which act as source for metal, are called Ore. From ore metal can be obtained economically .
3. The unwanted impurities present in ore are called Gangue.
4. The entire process of extraction of metal from its ore is called Metallurgy.
5. Removal of gangue from ore is called Concentration, Dressing or Benefaction of ore.
6. Concentration by Hydraulic washing is based on the difference in gravities of ore and gangue particles.