

**Question 1.1:**

Calculate the molecular mass of the following:

(i) H₂O (ii) CO₂ (iii) CH₄

Answer

(i) H₂O:

The molecular mass of water, H₂O

= (2 × Atomic mass of hydrogen) + (1 × Atomic mass of oxygen)

= [2(1.0084) + 1(16.00 u)]

= 2.016 u + 16.00 u

= 18.016

= 18.02 u

(ii) CO₂:

The molecular mass of carbon dioxide, CO₂

= (1 × Atomic mass of carbon) + (2 × Atomic mass of oxygen)

= [1(12.011 u) + 2 (16.00 u)]

= 12.011 u + 32.00 u

= 44.01 u

(iii) CH₄:

The molecular mass of methane, CH₄

= (1 × Atomic mass of carbon) + (4 × Atomic mass of hydrogen)

= [1(12.011 u) + 4 (1.008 u)]

= 12.011 u + 4.032 u

= 16.043 u

Question 1.2:

Calculate the mass percent of different elements present in sodium sulphate (Na₂SO₄).

Answer

The molecular formula of sodium sulphate is Na₂SO₄.

Molar mass of Na₂SO₄ = [(2 × 23.0) + (32.066) + 4 (16.00)]

= 142.066 g

Mass percent of an element = $\frac{\text{Mass of that element in the compound}}{\text{Molar mass of the compound}} \times 100$



∴ Mass percent of sodium:

$$\begin{aligned} &= \frac{46.0 \text{ g}}{142.066 \text{ g}} \times 100 \\ &= 32.379 \\ &= 32.4\% \end{aligned}$$

Mass percent of sulphur:

$$\begin{aligned} &= \frac{32.066 \text{ g}}{142.066 \text{ g}} \times 100 \\ &= 22.57 \\ &= 22.6\% \end{aligned}$$

Mass percent of oxygen:

$$\begin{aligned} &= \frac{64.0 \text{ g}}{142.066 \text{ g}} \times 100 \\ &= 45.049 \\ &= 45.05\% \end{aligned}$$

Question 1.3:

Determine the empirical formula of an oxide of iron which has 69.9% iron and 30.1% dioxygen by mass.

Answer

% of iron by mass = 69.9 % [Given]

% of oxygen by mass = 30.1 % [Given]

Relative moles of iron in iron oxide:

$$\begin{aligned} &= \frac{\% \text{ of iron by mass}}{\text{Atomic mass of iron}} \\ &= \frac{69.9}{55.85} \\ &= 1.25 \end{aligned}$$

Relative moles of oxygen in iron oxide:



$$\begin{aligned} &= \frac{\% \text{ of oxygen by mass}}{\text{Atomic mass of oxygen}} \\ &= \frac{30.1}{16.00} \\ &= 1.88 \end{aligned}$$

Simplest molar ratio of iron to oxygen:

$$= 1.25 : 1.88$$

$$= 1 : 1.5$$

$$\approx 2 : 3$$

∴ The empirical formula of the iron oxide is Fe_2O_3 .

Question 1.4:

Calculate the amount of carbon dioxide that could be produced when

- 1 mole of carbon is burnt in air.
- 1 mole of carbon is burnt in 16 g of dioxygen.
- 2 moles of carbon are burnt in 16 g of dioxygen.

Answer

The balanced reaction of combustion of carbon can be written as:

(i) As per the balanced equation, 1 mole of carbon burns in 1 mole of dioxygen (air) to produce 1 mole of carbon dioxide.

(ii) According to the question, only 16 g of dioxygen is available. Hence, it will react with 0.5 mole of carbon to give 22 g of carbon dioxide. Hence, it is a limiting reactant.

(iii) According to the question, only 16 g of dioxygen is available. It is a limiting reactant. Thus, 16 g of dioxygen can combine with only 0.5 mole of carbon to give 22 g of carbon dioxide.

Question 1.5:

Calculate the mass of sodium acetate (CH_3COONa) required to make 500 mL of 0.375 molar aqueous solution. Molar mass of sodium acetate is $82.0245 \text{ g mol}^{-1}$

Answer

0.375 M aqueous solution of sodium acetate

\equiv 1000 mL of solution containing 0.375 moles of sodium acetate

∴ Number of moles of sodium acetate in 500 mL



$$\begin{aligned} &= \frac{0.375}{1000} \times 500 \\ &= 0.1875 \text{ mole} \end{aligned}$$

Molar mass of sodium acetate = 82.0245 g mole⁻¹ (Given)

$$\begin{aligned} \therefore \text{Required mass of sodium acetate} &= (82.0245 \text{ g mol}^{-1}) (0.1875 \text{ mole}) \\ &= 15.38 \text{ g} \end{aligned}$$

Question 1.6:

Calculate the concentration of nitric acid in moles per litre in a sample which has a density, 1.41 g mL⁻¹ and the mass per cent of nitric acid in it being 69%.

Answer

Mass percent of nitric acid in the sample = 69 % [Given]

Thus, 100 g of nitric acid contains 69 g of nitric acid by mass.

Molar mass of nitric acid (HNO₃)

$$= \{1 + 14 + 3(16)\} \text{ g mol}^{-1}$$

$$= 1 + 14 + 48$$

$$= 63 \text{ g mol}^{-1}$$

∴ Number of moles in 69 g of HNO₃

$$= \frac{69\text{g}}{63 \text{ g mol}^{-1}}$$

$$= 1.095 \text{ mol}$$

Volume of 100g of nitric acid solution

$$= \frac{\text{Mass of solution}}{\text{density of solution}}$$

$$= \frac{100 \text{ g}}{1.41 \text{ g mL}^{-1}}$$

$$= 70.92 \text{ mL} \equiv 70.92 \times 10^{-3} \text{ L}$$

Concentration of nitric acid

$$= \frac{1.095 \text{ mole}}{70.92 \times 10^{-3} \text{ L}}$$

$$= 15.44 \text{ mol/L}$$

∴ Concentration of nitric acid = 15.44 mol/L

**Question 1.7:**

How much copper can be obtained from 100 g of copper sulphate (CuSO_4)?

Answer

1 mole of CuSO_4 contains 1 mole of copper.

Molar mass of $\text{CuSO}_4 = (63.5) + (32.00) + 4(16.00)$

$= 63.5 + 32.00 + 64.00$

$= 159.5 \text{ g}$

159.5 g of CuSO_4 contains 63.5 g of copper.

$\Rightarrow 100 \text{ g of } \text{CuSO}_4 \text{ will contain } \frac{63.5 \times 100 \text{ g}}{159.5} \text{ of copper.}$

\therefore Amount of copper that can be obtained from 100 g $\text{CuSO}_4 = \frac{63.5 \times 100}{159.5}$
 $= 39.81 \text{ g}$

Question 1.8:

Determine the molecular formula of an oxide of iron in which the mass per cent of iron and oxygen are 69.9 and 30.1 respectively. Given that the molar mass of the oxide is $159.69 \text{ g mol}^{-1}$.

Answer

Mass percent of iron (Fe) = 69.9% (Given)

Mass percent of oxygen (O) = 30.1% (Given)

Number of moles of iron present in the oxide $= \frac{69.90}{55.85}$
 $= 1.25$

Number of moles of oxygen present in the oxide $= \frac{30.1}{16.0}$
 $= 1.88$

Ratio of iron to oxygen in the oxide,

$= 1.25 : 1.88$

$= \frac{1.25}{1.25} : \frac{1.88}{1.25}$



$$= 1 : 1.5$$

$$= 2 : 3$$

∴ The empirical formula of the oxide is Fe_2O_3 .

$$\text{Empirical formula mass of } \text{Fe}_2\text{O}_3 = [2(55.85) + 3(16.00)] \text{ g}$$

$$\text{Molar mass of } \text{Fe}_2\text{O}_3 = 159.69 \text{ g}$$

$$\begin{aligned} \therefore n &= \frac{\text{Molar mass}}{\text{Empirical formula mass}} = \frac{159.69 \text{ g}}{159.7 \text{ g}} \\ &= 0.999 \\ &= 1(\text{approx}) \end{aligned}$$

Molecular formula of a compound is obtained by multiplying the empirical formula with n .

Thus, the empirical formula of the given oxide is Fe_2O_3 and n is 1.

Hence, the molecular formula of the oxide is Fe_2O_3 .

Question 1.9:

Calculate the atomic mass (average) of chlorine using the following data:

	% Natural Abundance	Molar Mass
^{35}Cl	75.77	34.9689
^{37}Cl	24.23	36.9659

Answer

The average atomic mass of chlorine

$$\begin{aligned} &= \left[\left(\frac{\text{Fractional abundance}}{\text{of } ^{35}\text{Cl}} \right) \left(\frac{\text{Molar mass}}{\text{of } ^{35}\text{Cl}} \right) + \left(\frac{\text{Fractional abundance}}{\text{of } ^{37}\text{Cl}} \right) \left(\frac{\text{Molar mass}}{\text{of } ^{37}\text{Cl}} \right) \right] \\ &= \left[\left\{ \left(\frac{75.77}{100} \right) (34.9689 \text{ u}) \right\} + \left\{ \left(\frac{24.23}{100} \right) (36.9659 \text{ u}) \right\} \right] \end{aligned}$$

$$= 26.4959 + 8.9568$$

$$= 35.4527 \text{ u}$$

∴ The average atomic mass of chlorine = 35.4527 u

Question 1.10:

In three moles of ethane (C_2H_6), calculate the following:



- (i) Number of moles of carbon atoms.
(ii) Number of moles of hydrogen atoms.
(iii) Number of molecules of ethane.

Answer

(i) 1 mole of C_2H_6 contains 2 moles of carbon atoms.

$$\begin{aligned} \therefore \text{Number of moles of carbon atoms in 3 moles of } C_2H_6 \\ = 2 \times 3 = 6 \end{aligned}$$

(ii) 1 mole of C_2H_6 contains 6 moles of hydrogen atoms.

$$\begin{aligned} \therefore \text{Number of moles of carbon atoms in 3 moles of } C_2H_6 \\ = 3 \times 6 = 18 \end{aligned}$$

(iii) 1 mole of C_2H_6 contains 6.023×10^{23} molecules of ethane.

$$\begin{aligned} \therefore \text{Number of molecules in 3 moles of } C_2H_6 \\ = 3 \times 6.023 \times 10^{23} = 18.069 \times 10^{23} \end{aligned}$$

Question 1.11:

What is the concentration of sugar ($C_{12}H_{22}O_{11}$) in mol L^{-1} if its 20 g are dissolved in enough water to make a final volume up to 2 L?

Answer

Molarity (M) of a solution is given by,

$$\begin{aligned} &= \frac{\text{Number of moles of solute}}{\text{Volume of solution in Litres}} \\ &= \frac{\text{Mass of sugar/molar mass of sugar}}{2 \text{ L}} \\ &= \frac{20\text{g} / [(12 \times 12) + (1 \times 22) + (11 \times 16)]\text{g}}{2 \text{ L}} \\ &= \frac{20\text{g} / 342 \text{ g}}{2 \text{ L}} \\ &= \frac{0.0585 \text{ mol}}{2 \text{ L}} \\ &= 0.02925 \text{ mol L}^{-1} \end{aligned}$$

$$\therefore \text{Molar concentration of sugar} = 0.02925 \text{ mol L}^{-1}$$

Question 1.12:



If the density of methanol is 0.793 kg L^{-1} , what is its volume needed for making 2.5 L of its 0.25 M solution?

Answer

Molar mass of methanol (CH_3OH) = $(1 \times 12) + (4 \times 1) + (1 \times 16)$

= 32 g mol^{-1}

= $0.032 \text{ kg mol}^{-1}$

Molarity of methanol solution = $\frac{0.793 \text{ kg L}^{-1}}{0.032 \text{ kg mol}^{-1}}$

= 24.78 mol L^{-1}

(Since density is mass per unit volume)

Applying,

$M_1V_1 = M_2V_2$

(Given solution) (Solution to be prepared)

$(24.78 \text{ mol L}^{-1}) V_1 = (2.5 \text{ L}) (0.25 \text{ mol L}^{-1})$

$V_1 = 0.0252 \text{ L}$

$V_1 = 25.22 \text{ mL}$

Question 1.13:

Pressure is determined as force per unit area of the surface. The SI unit of pressure, Pascal is as shown below:

$1 \text{ Pa} = 1 \text{ N m}^{-2}$

If mass of air at sea level is 1034 g cm^{-2} , calculate the pressure in Pascal.

Answer

Pressure is defined as force acting per unit area of the surface.

$$P = \frac{F}{A}$$
$$= \frac{1034 \text{ g} \times 9.8 \text{ ms}^{-2}}{\text{cm}^2} \times \frac{1 \text{ kg}}{1000 \text{ g}} \times \frac{(100)^2 \text{ cm}^2}{1 \text{ m}^2}$$

= $1.01332 \times 10^5 \text{ kg m}^{-1}\text{s}^{-2}$

We know,

$1 \text{ N} = 1 \text{ kg ms}^{-2}$

Then,



$$1 \text{ Pa} = 1 \text{ Nm}^{-2} = 1 \text{ kg m}^{-2}\text{s}^{-2}$$

$$1 \text{ Pa} = 1 \text{ kg m}^{-1}\text{s}^{-2}$$

$$\therefore \text{Pressure} = 1.01332 \times 10^5 \text{ Pa}$$

Question 1.14:

What is the SI unit of mass? How is it defined?

Answer

The SI unit of mass is kilogram (kg). 1 Kilogram is defined as the mass equal to the mass of the international prototype of kilogram.

Question 1.15:

Match the following prefixes with their multiples:

	Prefixes	Multiples
(i)	micro	10^6
(ii)	deca	10^9
(iii)	mega	10^{-6}
(iv)	giga	10^{-15}
(v)	femto	10

Answer

	Prefix	Multiples
(i)	micro	10^{-6}
(ii)	deca	10
(iii)	mega	10^6
(iv)	giga	10^9
(v)	femto	10^{-15}

**Question 1.16:**

What do you mean by significant figures?

Answer

Significant figures are those meaningful digits that are known with certainty.

They indicate uncertainty in an experiment or calculated value. For example, if 15.6 mL is the result of an experiment, then 15 is certain while 6 is uncertain, and the total number of significant figures are 3.

Hence, significant figures are defined as the total number of digits in a number including the last digit that represents the uncertainty of the result.

Question 1.17:

A sample of drinking water was found to be severely contaminated with chloroform, CHCl_3 , supposed to be carcinogenic in nature. The level of contamination was 15 ppm (by mass).

(i) Express this in percent by mass.

(ii) Determine the molality of chloroform in the water sample.

Answer

(i) 1 ppm is equivalent to 1 part out of 1 million (10^6) parts.

∴ Mass percent of 15 ppm chloroform in water

$$\begin{aligned} &= \frac{15}{10^6} \times 100 \\ &\approx 1.5 \times 10^{-3} \% \end{aligned}$$

(ii) 100 g of the sample contains 1.5×10^{-3} g of CHCl_3 .

⇒ 1000 g of the sample contains 1.5×10^{-2} g of CHCl_3 .

∴ Molality of chloroform in water

$$= \frac{1.5 \times 10^{-2} \text{ g}}{\text{Molar mass of } \text{CHCl}_3}$$

Molar mass of $\text{CHCl}_3 = 12.00 + 1.00 + 3(35.5)$

$= 119.5 \text{ g mol}^{-1}$

∴ Molality of chloroform in water $= 0.0125 \times 10^{-2} \text{ m}$

$= 1.25 \times 10^{-4} \text{ m}$

**Question 1.18:**

Express the following in the scientific notation:

- (i) 0.0048
- (ii) 234,000
- (iii) 8008
- (iv) 500.0
- (v) 6.0012

Answer

- (i)** $0.0048 = 4.8 \times 10^{-3}$
- (ii)** $234,000 = 2.34 \times 10^5$
- (iii)** $8008 = 8.008 \times 10^3$
- (iv)** $500.0 = 5.000 \times 10^2$
- (v)** $6.0012 = 6.0012$

Question 1.19:

How many significant figures are present in the following?

- (i) 0.0025
- (ii) 208
- (iii) 5005
- (iv) 126,000
- (v) 500.0
- (vi) 2.0034

Answer

- (i)** 0.0025

There are 2 significant figures.

- (ii)** 208

There are 3 significant figures.

- (iii)** 5005

There are 4 significant figures.

- (iv)** 126,000

There are 3 significant figures.



(v) 500.0

There are 4 significant figures.

(vi) 2.0034

There are 5 significant figures.

Question 1.20:

Round up the following upto three significant figures:

(i) 34.216

(ii) 10.4107

(iii) 0.04597

(iv) 2808

Answer

(i) 34.2

(ii) 10.4

(iii) 0.0460

(iv) 2810

Question 1.21:

The following data are obtained when dinitrogen and dioxygen react together to form different compounds:

	Mass of dinitrogen	Mass of dioxygen
(i)	14 g	16 g
(ii)	14 g	32 g
(iii)	28 g	32 g
(iv)	28 g	80 g

(a) Which law of chemical combination is obeyed by the above experimental data? Give its statement.

(b) Fill in the blanks in the following conversions:

(i) 1 km = mm = pm

(ii) 1 mg = kg = ng

(iii) 1 mL = L = dm³

Answer

(a)



If we fix the mass of dinitrogen at 28 g, then the masses of dioxygen that will combine with the fixed mass of dinitrogen are 32 g, 64 g, 32 g, and 80 g.

The masses of dioxygen bear a whole number ratio of 1:2:2:5. Hence, the given experimental data obeys the law of multiple proportions. The law states that if two elements combine to form more than one compound, then the masses of one element that combines with the fixed mass of another element are in the ratio of small whole numbers.

(b)

$$(i) 1 \text{ km} = 1 \text{ km} \times \frac{1000 \text{ m}}{1 \text{ km}} \times \frac{100 \text{ cm}}{1 \text{ m}} \times \frac{10 \text{ mm}}{1 \text{ cm}}$$
$$\therefore 1 \text{ km} = 10^6 \text{ mm}$$

$$1 \text{ km} = 1 \text{ km} \times \frac{1000 \text{ m}}{1 \text{ km}} \times \frac{1 \text{ pm}}{10^{-12} \text{ m}}$$
$$\therefore 1 \text{ km} = 10^{15} \text{ pm}$$

$$\text{Hence, } 1 \text{ km} = 10^6 \text{ mm} = 10^{15} \text{ pm}$$

$$(ii) 1 \text{ mg} = 1 \text{ mg} \times \frac{1 \text{ g}}{1000 \text{ mg}} \times \frac{1 \text{ kg}}{1000 \text{ g}}$$
$$\Rightarrow 1 \text{ mg} = 10^{-6} \text{ kg}$$

$$1 \text{ mg} = 1 \text{ mg} \times \frac{1 \text{ g}}{1000 \text{ mg}} \times \frac{1 \text{ ng}}{10^{-9} \text{ g}}$$
$$\Rightarrow 1 \text{ mg} = 10^6 \text{ ng}$$
$$\therefore 1 \text{ mg} = 10^{-6} \text{ kg} = 10^6 \text{ ng}$$

$$(iii) 1 \text{ mL} = 1 \text{ mL} \times \frac{1 \text{ L}}{1000 \text{ mL}}$$
$$\Rightarrow 1 \text{ mL} = 10^{-3} \text{ L}$$

$$1 \text{ mL} = 1 \text{ cm}^3 = 1 \text{ cm}^3 \frac{1 \text{ dm} \times 1 \text{ dm} \times 1 \text{ dm}}{10 \text{ cm} \times 10 \text{ cm} \times 10 \text{ cm}}$$
$$\Rightarrow 1 \text{ mL} = 10^{-3} \text{ dm}^3$$
$$\therefore 1 \text{ mL} = 10^{-3} \text{ L} = 10^{-3} \text{ dm}^3$$

**Question 1.22:**

If the speed of light is $3.0 \times 10^8 \text{ m s}^{-1}$, calculate the distance covered by light in 2.00 ns.

Answer

According to the question:

Time taken to cover the distance = 2.00 ns

= $2.00 \times 10^{-9} \text{ s}$

Speed of light = $3.0 \times 10^8 \text{ ms}^{-1}$

Distance travelled by light in 2.00 ns

= Speed of light \times Time taken


= $(3.0 \times 10^8 \text{ ms}^{-1}) (2.00 \times 10^{-9} \text{ s})$

= $6.00 \times 10^{-1} \text{ m}$

= 0.600 m

Question 1.23:

In a reaction



Identify the limiting reagent, if any, in the following reaction mixtures.

(i) 300 atoms of A + 200 molecules of B

(ii) 2 mol A + 3 mol B

(iii) 100 atoms of A + 100 molecules of B

(iv) 5 mol A + 2.5 mol B

(v) 2.5 mol A + 5 mol B

Answer

A limiting reagent determines the extent of a reaction. It is the reactant which is the first to get consumed during a reaction, thereby causing the reaction to stop and limiting the amount of products formed.

(i) According to the given reaction, 1 atom of A reacts with 1 molecule of B. Thus, 200 molecules of B will react with 200 atoms of A, thereby leaving 100 atoms of A unused. Hence, B is the limiting reagent.

(ii) According to the reaction, 1 mol of A reacts with 1 mol of B. Thus, 2 mol of A will react with only 2 mol of B. As a result, 1 mol of A will not be consumed. Hence, A is the limiting reagent.




(iii) According to the given reaction, 1 atom of A combines with 1 molecule of B. Thus, all 100 atoms of A will combine with all 100 molecules of B. Hence, the mixture is stoichiometric where no limiting reagent is present.

(iv) 1 mol of atom A combines with 1 mol of molecule B. Thus, 2.5 mol of B will combine with only 2.5 mol of A. As a result, 2.5 mol of A will be left as such. Hence, B is the limiting reagent.

(v) According to the reaction, 1 mol of atom A combines with 1 mol of molecule B. Thus, 2.5 mol of A will combine with only 2.5 mol of B and the remaining 2.5 mol of B will be left as such. Hence, A is the limiting reagent.

Question 1.24:

Dinitrogen and dihydrogen react with each other to produce ammonia according to the following chemical equation:




(i) Calculate the mass of ammonia produced if 2.00×10^3 g dinitrogen reacts with 1.00×10^3 g of dihydrogen.

(ii) Will any of the two reactants remain unreacted?

(iii) If yes, which one and what would be its mass?

Answer

(i) Balancing the given chemical equation,



From the equation, 1 mole (28 g) of dinitrogen reacts with 3 mole (6 g) of dihydrogen to give 2 mole (34 g) of ammonia.

$\Rightarrow 2.00 \times 10^3$ g of dinitrogen will react with $\frac{6 \text{ g}}{28 \text{ g}} \times 2.00 \times 10^3 \text{ g}$ dihydrogen i.e.,

2.00×10^3 g of dinitrogen will react with 428.6 g of dihydrogen.

Given,

Amount of dihydrogen = 1.00×10^3 g

Hence, N_2 is the limiting reagent.

\therefore 28 g of N_2 produces 34 g of NH_3 .

Hence, mass of ammonia produced by 2000 g of N_2 = $\frac{34 \text{ g}}{28 \text{ g}} \times 2000 \text{ g}$



= 2428.57 g

(ii) N_2 is the limiting reagent and H_2 is the excess reagent. Hence, H_2 will remain unreacted.

(iii) Mass of dihydrogen left unreacted = $1.00 \times 10^3 \text{ g} - 428.6 \text{ g}$
= 571.4 g

Question 1.25:

How are 0.50 mol Na_2CO_3 and 0.50 M Na_2CO_3 different?

Answer

Molar mass of Na_2CO_3 = $(2 \times 23) + 12.00 + (3 \times 16)$
= 106 g mol^{-1}

Now, 1 mole of Na_2CO_3 means 106 g of Na_2CO_3 .

$\therefore 0.5 \text{ mol of } Na_2CO_3 = \frac{106 \text{ g}}{1 \text{ mole}} \times 0.5 \text{ mol } Na_2CO_3$
= 53 g Na_2CO_3

$\Rightarrow 0.50 \text{ M of } Na_2CO_3 = 0.50 \text{ mol/L } Na_2CO_3$


Hence, 0.50 mol of Na_2CO_3 is present in 1 L of water or 53 g of Na_2CO_3 is present in 1 L of water.

Question 1.26:

If ten volumes of dihydrogen gas react with five volumes of dioxygen gas, how many volumes of water vapour would be produced?

Answer

Reaction of dihydrogen with dioxygen can be written as:



Now, two volumes of dihydrogen react with one volume of dihydrogen to produce two volumes of water vapour.

Hence, ten volumes of dihydrogen will react with five volumes of dioxygen to produce ten volumes of water vapour.

Question 1.27:

Convert the following into basic units:



- (i) 28.7 pm
(ii) 15.15 pm
(iii) 25365 mg

Answer

(i) 28.7 pm:

$$1 \text{ pm} = 10^{-12} \text{ m}$$

$$\begin{aligned} \therefore 28.7 \text{ pm} &= 28.7 \times 10^{-12} \text{ m} \\ &= 2.87 \times 10^{-11} \text{ m} \end{aligned}$$

(ii) 15.15 pm:

$$1 \text{ pm} = 10^{-12} \text{ m}$$

$$\begin{aligned} \therefore 15.15 \text{ pm} &= 15.15 \times 10^{-12} \text{ m} \\ &= 1.515 \times 10^{-11} \text{ m} \end{aligned}$$

(iii) 25365 mg:

$$1 \text{ mg} = 10^{-3} \text{ g}$$

$$25365 \text{ mg} = 2.5365 \times 10^4 \times 10^{-3} \text{ g}$$

Since,

$$1 \text{ g} = 10^{-3} \text{ kg}$$

$$2.5365 \times 10^1 \text{ g} = 2.5365 \times 10^{-1} \times 10^{-3} \text{ kg}$$

$$\therefore 25365 \text{ mg} = 2.5365 \times 10^{-2} \text{ kg}$$

Question 1.28:

Which one of the following will have largest number of atoms?

- (i) 1 g Au (s)
(ii) 1 g Na (s)
(iii) 1 g Li (s)
(iv) 1 g of Cl_2 (g)

Answer

$$\begin{aligned} 1 \text{ g of Au (s)} &= \frac{1}{197} \text{ mol of Au (s)} \\ &= \frac{6.022 \times 10^{23}}{197} \text{ atoms of Au (s)} \\ &= 3.06 \times 10^{21} \text{ atoms of Au (s)} \end{aligned}$$



$$\begin{aligned}1 \text{ g of Na (s)} &= \frac{1}{23} \text{ mol of Na (s)} \\ &= \frac{6.022 \times 10^{23}}{23} \text{ atoms of Na (s)} \\ &= 0.262 \times 10^{23} \text{ atoms of Na (s)} \\ &= 26.2 \times 10^{21} \text{ atoms of Na (s)}\end{aligned}$$

$$\begin{aligned}1 \text{ g of Li (s)} &= \frac{1}{7} \text{ mol of Li (s)} \\ &= \frac{6.022 \times 10^{23}}{7} \text{ atoms of Li (s)} \\ &= 0.86 \times 10^{23} \text{ atoms of Li (s)} \\ &= 86.0 \times 10^{21} \text{ atoms of Li (s)}\end{aligned}$$

$$\begin{aligned}1 \text{ g of Cl}_2 \text{ (g)} &= \frac{1}{71} \text{ mol of Cl}_2 \text{ (g)} \\ (\text{Molar mass of Cl}_2 \text{ molecule} &= 35.5 \times 2 = 71 \text{ g mol}^{-1})\end{aligned}$$

$$\begin{aligned}&= \frac{6.022 \times 10^{23}}{71} \text{ atoms of Cl}_2 \text{ (g)} \\ &= 0.0848 \times 10^{23} \text{ atoms of Cl}_2 \text{ (g)} \\ &= 8.48 \times 10^{21} \text{ atoms of Cl}_2 \text{ (g)}\end{aligned}$$

Hence, 1 g of Li (s) will have the largest number of atoms.

Question 1.29:

Calculate the molarity of a solution of ethanol in water in which the mole fraction of ethanol is 0.040 (assume the density of water to be one).

Answer

$$\text{Mole fraction of C}_2\text{H}_5\text{OH} = \frac{\text{Number of moles of C}_2\text{H}_5\text{OH}}{\text{Number of moles of solution}}$$

$$0.040 = \frac{n_{\text{C}_2\text{H}_5\text{OH}}}{n_{\text{C}_2\text{H}_5\text{OH}} + n_{\text{H}_2\text{O}}} \dots\dots\dots(1)$$

Number of moles present in 1 L water:



$$n_{\text{H}_2\text{O}} = \frac{1000 \text{ g}}{18 \text{ g mol}^{-1}}$$

$$n_{\text{H}_2\text{O}} = 55.55 \text{ mol}$$

Substituting the value of $n_{\text{H}_2\text{O}}$ in equation (1),

$$\frac{n_{\text{C}_2\text{H}_5\text{OH}}}{n_{\text{C}_2\text{H}_5\text{OH}} + 55.55} = 0.040$$

$$n_{\text{C}_2\text{H}_5\text{OH}} = 0.040n_{\text{C}_2\text{H}_5\text{OH}} + (0.040)(55.55)$$

$$0.96n_{\text{C}_2\text{H}_5\text{OH}} = 2.222 \text{ mol}$$

$$n_{\text{C}_2\text{H}_5\text{OH}} = \frac{2.222}{0.96} \text{ mol}$$

$$n_{\text{C}_2\text{H}_5\text{OH}} = 2.314 \text{ mol}$$

$$\therefore \text{Molarity of solution} = \frac{2.314 \text{ mol}}{1 \text{ L}}$$

$$= 2.314 \text{ M}$$

Question 1.30:

What will be the mass of one ^{12}C atom in g?

Answer

1 mole of carbon atoms = 6.023×10^{23} atoms of carbon

= 12 g of carbon

$$\therefore \text{Mass of one } ^{12}\text{C} \text{ atom} = \frac{12 \text{ g}}{6.022 \times 10^{23}}$$

$$= 1.993 \times 10^{-23} \text{ g}$$

Question 1.31:

How many significant figures should be present in the answer of the following calculations?

$$(i) \frac{0.02856 \times 298.15 \times 0.112}{0.5785}$$

$$(ii) 5 \times 5.364$$

$$(iii) 0.0125 + 0.7864 + 0.0215$$



Answer

$$(i) \frac{0.02856 \times 298.15 \times 0.112}{0.5785}$$

Least precise number of calculation = 0.112

∴ Number of significant figures in the answer

= Number of significant figures in the least precise number

= 3

$$(ii) 5 \times 5.364$$

Least precise number of calculation = 5.364

∴ Number of significant figures in the answer = Number of significant figures in 5.364

= 4

$$(iii) 0.0125 + 0.7864 + 0.0215$$

Since the least number of decimal places in each term is four, the number of significant figures in the answer is also 4.

Question 1.32:

Use the data given in the following table to calculate the molar mass of naturally occurring *argon* isotopes:

Isotope	Isotopic molar mass	Abundance
^{36}Ar	35.96755 gmol^{-1}	0.337%
^{38}Ar	37.96272 gmol^{-1}	0.063%
^{40}Ar	39.9624 gmol^{-1}	99.600%

Answer

Molar mass of argon

$$\begin{aligned} &= \left[\left(35.96755 \times \frac{0.337}{100} \right) + \left(37.96272 \times \frac{0.063}{100} \right) + \left(39.9624 \times \frac{99.60}{100} \right) \right] \text{gmol}^{-1} \\ &= [0.121 + 0.024 + 39.802] \text{gmol}^{-1} \\ &= 39.947 \text{gmol}^{-1} \end{aligned}$$

Question 1.33:



Calculate the number of atoms in each of the following (i) 52 moles of Ar (ii) 52 u of He (iii) 52 g of He.

Answer

(i) 1 mole of Ar = 6.022×10^{23} atoms of Ar

$$\begin{aligned} \therefore 52 \text{ mol of Ar} &= 52 \times 6.022 \times 10^{23} \text{ atoms of Ar} \\ &= 3.131 \times 10^{25} \text{ atoms of Ar} \end{aligned}$$

(ii) 1 atom of He = 4 u of He

Or,

4 u of He = 1 atom of He

$$1 \text{ u of He} = \frac{1}{4} \text{ atom of He}$$

$$52 \text{ u of He} = \frac{52}{4} \text{ atom of He}$$

= 13 atoms of He

(iii) 4 g of He = 6.022×10^{23} atoms of He

$$\begin{aligned} \therefore 52 \text{ g of He} &= \frac{6.022 \times 10^{23} \times 52}{4} \text{ atoms of He} \\ &= 7.8286 \times 10^{24} \text{ atoms of He} \end{aligned}$$

Question 1.34:

A welding fuel gas contains carbon and hydrogen only. Burning a small sample of it in oxygen gives 3.38 g carbon dioxide, 0.690 g of water and no other products. A volume of 10.0 L (measured at STP) of this welding gas is found to weigh 11.6 g. Calculate (i) empirical formula, (ii) molar mass of the gas, and (iii) molecular formula.

Answer

(i) 1 mole (44 g) of CO_2 contains 12 g of carbon.

$$\begin{aligned} \therefore 3.38 \text{ g of } \text{CO}_2 \text{ will contain carbon} &= \frac{12 \text{ g}}{44 \text{ g}} \times 3.38 \text{ g} \\ &= 0.9217 \text{ g} \end{aligned}$$

18 g of water contains 2 g of hydrogen.

$$\therefore 0.690 \text{ g of water will contain hydrogen} = \frac{2 \text{ g}}{18 \text{ g}} \times 0.690$$



$$= 0.0767 \text{ g}$$

Since carbon and hydrogen are the only constituents of the compound, the total mass of the compound is:

$$= 0.9217 \text{ g} + 0.0767 \text{ g}$$

$$= 0.9984 \text{ g}$$

$$\therefore \text{Percent of C in the compound} = \frac{0.9217 \text{ g}}{0.9984 \text{ g}} \times 100$$

$$= 92.32\%$$

$$\text{Percent of H in the compound} = \frac{0.0767 \text{ g}}{0.9984 \text{ g}} \times 100$$

$$= 7.68\%$$

$$\text{Moles of carbon in the compound} = \frac{92.32}{12.00}$$

$$= 7.69$$

$$\text{Moles of hydrogen in the compound} = \frac{7.68}{1}$$

$$= 7.68$$

$$\therefore \text{Ratio of carbon to hydrogen in the compound} = 7.69 : 7.68$$

$$= 1 : 1$$

Hence, the empirical formula of the gas is CH.

(ii) Given,

$$\text{Weight of 10.0L of the gas (at S.T.P)} = 11.6 \text{ g}$$

$$\therefore \text{Weight of 22.4 L of gas at STP} = \frac{11.6 \text{ g}}{10.0\text{L}} \times 22.4 \text{ L}$$

$$= 25.984 \text{ g}$$

$$\approx 26 \text{ g}$$

Hence, the molar mass of the gas is 26 g.

(iii) Empirical formula mass of CH = 12 + 1 = 13 g

$$n = \frac{\text{Molar mass of gas}}{\text{Empirical formula mass of gas}}$$
$$= \frac{26 \text{ g}}{13 \text{ g}}$$



$$n = 2$$

$$\begin{aligned} \therefore \text{Molecular formula of gas} &= (\text{CH})_n \\ &= \text{C}_2\text{H}_2 \end{aligned}$$

Question 1.35:

Calcium carbonate reacts with aqueous HCl to give CaCl_2 and CO_2 according to the reaction, $\text{CaCO}_{3(s)} + 2\text{HCl}_{(aq)} \rightarrow \text{CaCl}_{2(aq)} + \text{CO}_{2(g)} + \text{H}_2\text{O}_{(l)}$

What mass of CaCO_3 is required to react completely with 25 mL of 0.75 M HCl?

Answer

$$\begin{aligned} 0.75 \text{ M of HCl} &\equiv 0.75 \text{ mol of HCl are present in 1 L of water} \\ &\equiv [(0.75 \text{ mol}) \times (36.5 \text{ g mol}^{-1})] \text{ HCl is present in 1 L of water} \\ &\equiv 27.375 \text{ g of HCl is present in 1 L of water} \end{aligned}$$


Thus, 1000 mL of solution contains 27.375 g of HCl.

\therefore Amount of HCl present in 25 mL of solution

$$= \frac{27.375 \text{ g}}{1000 \text{ mL}} \times 25 \text{ mL}$$

$$= 0.6844 \text{ g}$$

From the given chemical equation,




2 mol of HCl ($2 \times 36.5 = 71 \text{ g}$) react with 1 mol of CaCO_3 (100 g).

$$\therefore \text{Amount of } \text{CaCO}_3 \text{ that will react with } 0.6844 \text{ g} = \frac{100}{71} \times 0.6844 \text{ g}$$

$$= 0.9639 \text{ g}$$

Question 1.36:

Chlorine is prepared in the laboratory by treating manganese dioxide (MnO_2) with aqueous hydrochloric acid according to the reaction



How many grams of HCl react with 5.0 g of manganese dioxide?

Answer

1 mol [$55 + 2 \times 16 = 87 \text{ g}$] MnO_2 reacts completely with 4 mol [$4 \times 36.5 = 146 \text{ g}$] of HCl.



∴ 5.0 g of MnO_2 will react with

$$= \frac{146 \text{ g}}{87 \text{ g}} \times 5.0 \text{ g} \text{ of HCl}$$

$$= 8.4 \text{ g of HCl}$$

Hence, 8.4 g of HCl will react completely with 5.0 g of manganese dioxide.

UNIT-1

SOME BASIC CONCEPTS OF CHEMISTRY

Law of conservation of mass : ‘Mass can neither be created nor destroyed.’
In all physical and chemical changes, the total mass of reactants is equal to that of products.

Law of constant composition : A chemical compound is always found to be made of same elements combined together in the same fixed ratio by mass.

Law of multiple proportion : Two elements combined together to form two or more chemical compounds then the masses of the elements which combine with a fixed mass of another bear a simple ratio to one another.

Gram atomic mass or molar mass of an element is mass of 1 mol of atoms or atomic mass expressed in grams. For example, atomic mass of Ag = 108 u, therefore, molar mass of Ag is 108 grams per mol. Molar mass of an element is also called one gram atom.

Gram molecular mass or the molar mass of molecular substances is the mass of 1 mol of molecules or molecular mass expressed in grams. For example, molecular mass of CO₂ is 44 u, therefore, molar mass of CO₂ is 44 grams/mol.

Molar mass of ionic substance is the mass of 1 mol of formula units of ionic substance.

Molar mass and standard molar volume of gaseous substances :

1 mole of any gas occupies a volume of 22.4 L at STP, *i.e.*, at 298 K and 1 atm. If standard pressure is taken as 1 bar, then the standard molar volume is taken as 22.7 L.

$$\text{Molarity (M)} = \frac{W_B \times 1000}{M_B \times V_{\text{mL}}}$$
$$\text{Molality (m)} = \frac{W_B \times 1000}{M_B \times W_A}$$

For binary solutions : Mole fraction (X_B) of solute = $\frac{n_B}{n_A + n_B}$
 $X_A = 1 - X_B$

where X_A = mole fraction of solvent,
 X_B = mol fraction of solute
 W_A = mass of solvent
 W_B = Mass of solute
 M_B = Molar mass of solute
 V_{mL} = Volume of solution in mL

Dilution Formula : $M_1V_1 = M_2V_2$

Where : M_1 = Molarity of concentrated solution

V_1 = Volume of concentrated solution

M_2 = Molarity of dilute solution


V_2 = Volume of dilute solution

For a general chemical equation : $aA + bB \rightarrow cC + dD$

Molarity relationship is : mol of B = mol of A $\times \frac{b \text{ mol of B}}{a \text{ mol of A}}$

Example : Calculate the volume of 0.1 M NaOH solution required to neutralise the solution produced by dissolving 1.1 g of P_4O_6 in water.

Solution :



$$\text{mol of } P_4O_6 = 1.1 \text{ g } P_4O_6 \times \frac{1 \text{ mol } P_4O_6}{220 \text{ g } P_4O_6} = 0.5 \times 10^{-2} \text{ mol } P_4O_6$$

$$\begin{aligned} \text{mol of NaOH} &= 0.5 \times 10^{-2} \text{ mol } P_4O_6 \times \frac{4 \text{ mol } H_3PO_3}{1 \text{ mol } P_4O_6} \times \frac{2 \text{ mol NaOH}}{1 \text{ mol } H_3PO_3} \\ &= 4 \times 10^{-2} \text{ mol NaOH} \end{aligned}$$

$$\begin{aligned} \therefore \text{Volume of NaOH solution in litres} &= 4 \times 10^{-2} \text{ mol NaOH} \times \frac{1 \text{ L NaOH solution}}{0.1 \text{ mol NaOH}} \\ &= 4 \times 10^{-1} \text{ L} = 0.4 \text{ L} \end{aligned}$$

1- MARK QUESTIONS

1. Why can't solids be compressed ?
2. Liquids take the shape of the container in which they are placed. Why ?
3. Give two examples of a homogeneous mixture.
4. Calculate the number of molecules present in 100.0 g of water.
5. Calculate the number of moles of oxygen atoms present in 22.0 g CO_2 .
6. Write the SI unit of temperature.
7. Define law of multiple proportions.

8. N_2 and H_2 combine according to the following equation :



If 100 mL of N_2 gas combines with 300 mL of H_2 gas, calculate the volume of NH_3 produced at same temperature and pressure.

9. Out of 1 M NaCl solution and 1 m NaCl solution, which one is more concentrated ?
10. Write the S.I. unit of Avogadro constant.
11. How much CO_2 is produced when 6.0 g C is burnt in excess oxygen ?
12. Molarity is temperature dependent but molality is not. Why?
13. How many moles of HCl are present in 100 ml of 12 M HCl solution ?
Ans : 1.2 mol
14. Calculate the mole fraction of N_2 when 28 g N_2 is mixed with 64 g O_2 gas.
Ans. : 0.50
15. A water sample has 20.0 ppm (by mass) Cl_2 present in it. Calculate the quantity of Cl_2 present in 100 ml water. (Density of water = 1.0 g ml^{-1} .)
16. Calculate the number of electrons in 17.0 g NH_3 .
17. Calculate the number of atoms present in 64.0 u helium.
18. If the density of 68% nitric acid solution is 1.41 g mL, calculate the mass of HNO_3 present in 1.0 L solution.
19. Express the strength of 0.5 M Na_2CO_3 solution in grams per litre.
20. Balance the following equation :
 $Mg + N_2 \rightarrow Mg_3N_2$
21. Calculate the number of g-atoms of sulphur (S_8) in 8.0 g sample.
22. Calculate the mass of NaOH required to make 250 mL of $\frac{M}{20}$ solution.
23. Which of the following has highest mass ?
(a) 32 g O_2 gas (b) 2 g atom of Cl_2
(c) 0.5 mol Fe (d) 9.03×10^{23} atoms of C
24. How many moles of methane are required to produce 22.0 g CO_2 in combustion reaction ?

2 - MARK QUESTIONS

1. Calculate the mass percentage of Na and Cl atoms in common salt. (Given molar mass NaCl = 58.5 g/mol)
2. How many significant figures are present in ?
(a) 126000 (b) 126.0
3. How much CO_2 is produced when 1.0 mol of carbon is burnt in 16.0 g oxygen ?
4. 0.5 mol each of H_2S and SO_2 are mixed together in a reaction flask in which the following reaction takes place :
 $2H_2S(g) + SO_2(g) \rightarrow 2H_2O(l) + 3S(s)$
Calculate the number of moles of sulphur formed.

5. Pure oxygen is prepared by thermal decomposition of KClO_3 according to the equation :



Calculate the volume of oxygen gas liberated at STP by heating 12.25 g $\text{KClO}_3(s)$.

6. Classify the following as pure substance or mixture ?
 (a) Ethyl alcohol (b) Blood
 (c) 22 carat gold (d) Air
7. How many significant figures are present in the answer of following calculations :
 (a) $0.0125 + 0.7864 - 0.023$ (b) $\frac{0.025 \times 298.15 \times 0.1155}{0.5785}$
8. Which of the following samples has the largest number of atoms :
 (a) 1.0 g $\text{H}_2(g)$ (b) 1.0 g $\text{Na}(s)$
 (c) 1.0 g $\text{CH}_3\text{OH}(l)$ (d) 1.0 g $\text{Br}_2(l)$
9. Determine the molecular formula of an oxide of iron in which the mass percent of iron and oxygen are 69.9 and 30.1 respectively. Molar mass of this oxide is 170.0 g mol^{-1} . [Ans. : Fe_2O_3]
10. The density of 3 M solution of NaCl is 1.25 g mL^{-1} . Calculate molality of solution. (Molar mass of NaCl is 58.5 g mol^{-1}).
11. Calculate the molarity of an aqueous solution of methanol in which the mole fraction of ethanol is 0.040. Assume the density of water to be 1.0 g mL^{-1} . [Ans. : 2.31 M]
12. How many grams of HCl react with 5.0 g MnO_2 according to the equation :
 $4\text{HCl}(aq) + \text{MnO}_2(s) \rightarrow 2\text{H}_2\text{O}(l) + \text{MnCl}_2(aq) + \text{Cl}_2(g)$
 [Ans. : 8.40 g]
13. How are 0.5 mol Na_2CO_3 and 0.5 M Na_2CO_3 are different from each other?
14. If mass of air at sea level is 1034 g cm^{-2} , calculate the pressure in pascal. (Given $g = 9.8 \text{ ms}^{-2}$, $1 \text{ Pa} = 1 \text{ Nm}^{-2}$) [Ans. : $1.01332 \times 10^5 \text{ Pa}$]
15. A polluted water sample has been found (CHCl_3) to have 15 ppm CHCl_3 in it.
 (a) Express this value in percent by mass.
 (b) Determine the molality of chloroform (CHCl_3) in the water sample.
 Ans. : (a) $\sim 15 \times 10^{-4} \text{ g}$ (b) $= 1.25 \times 10^{-4} \text{ g}$
16. Use the following data to calculate the molar mass of naturally occurring argon :

Isotope	Isotopic molar mass	Abundance
^{36}Ar	$35.96755 \text{ g mol}^{-1}$	0.337%
^{38}Ar	$37.96272 \text{ g mol}^{-1}$	0.063%
^{40}Ar	$35.9624 \text{ g mol}^{-1}$	99.600%

[Ans.: 39.948 g/mol]

17. If the speed of light is $3.0 \times 10^8 \text{ ms}^{-1}$, calculate the distance covered by light in 2.00 ns. [Ans.: 0.600 m]
18. State the law of definite proportions. Explain it with the help of an example.
19. Burning a sample of a hydrocarbon gas gives 3.38 g CO_2 and 0.690 g H_2O . A volume of 10L (measured at STP) of this hydrocarbon weighs 11.6 g. Calculate the molecular formula of this hydrocarbon. [Ans.: $\text{C}_2 \text{H}_2$]
20. In three moles of ethane (C_2H_6), Calculate the following :
- Number of moles of carbon atoms
 - Number of moles of Hydrogen atoms
 - Number of Molecules of ethane

3 - MARK QUESTIONS

- State and explain Avogadro's law. Illustrate it with an example.
 - 10.0 L of a welding gas weighs 11.6 g at STP. Calculate the molar mass of this gas. [Ans.: 26.0 g/mol]
- Calculate the mass of CaCO_3 required to react completely with 25mL of 0.75 M HCl. [Ans.: 0.938 g]
 - Calculate volume of CO_2 released at STP in this reaction. [Ans.: 0.21 L]
- Dinitrogen and dihydrogen react with each other to produce ammonia according to following chemical equation :

$$\text{N}_2 (\text{g}) + 3\text{H}_2 (\text{g}) \rightarrow 2\text{NH}_3 (\text{g})$$
 - Calculate the mass of ammonia gas formed if 2.0 kg of nitrogen gas reacts with 1.0 kg of hydrogen gas.
 - Which of the two reactants is the limiting reagent and why ?
 - Which of the two reactants will remain unreacted and what will be the amount left unreacted ? [Ans.: m (NH_3) = 2.571 kg, H_2 will remain unreacted its mass is 571.5 g]
- Calculate the molarity of solution prepared by dissolving 175.5 g NaCl in enough water to form 1.0 L of brine solution.
 - Calculate molality of solution if its density is 1.25 g ml^{-1} .
 - Calculate the mole fraction of NaCl.
- Calculate the number of atoms in :
 - 5.0 L oxygen gas at STP
 - 4.4 g of CO_2
 - 52 u of He

6. Calculate the number of moles :
- (a) 05. L of 0.75 M Na_2CO_3 (b) 7.85 g iron
- (c) 34.2 g sucrose ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$)
7. A compound contains 4.07% hydrogen, 24.27% carbon and 71.65% chlorine. Its molar mass is 98.96 g. Determine its empirical and molecular formulas.
8. In a reaction $\text{A} + \text{B}_2 \longrightarrow \text{AB}_2$
Identify the limiting reagent, if any in the following reaction mixtures
- (i) 300 atom of A + 200 molecules of B_2
- (ii) 2 mole of A + 3 mol of B_2
- (iii) 100 atom of A + 100 molecules of B_2
9. Calculate the mass of Na which contain the same number of atom as are present in 4 gram of calcium

5 - MARK QUESTIONS

1. (a) Write the difference between a homogenous and a heterogeneous mixture.
- (b) State Gay Lussac's Law of gaseous volumes.
- (c) Calculate the volume of 0.1 M NaOH solution is required to neutralise 100 mL of concentrated aqueous sulphuric acid which contains 98% H_2SO_4 by mass. The density of conc. H_2SO_4 is 1.84 g/mL.
NaOH reacts with H_2SO_4 according to the following equation :
- $$2\text{NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$$
- (At. mass/g mol⁻¹ H = 1, S = 32, O = 16)
2. (a) How much Cu can be obtained from 100 g CuSO_4 ?
- (b) Boron occurs in nature in the form of two isotopes $^{11}_5\text{B}$ and $^{10}_5\text{B}$ in ratio 81% and 19% respectively. Calculate its average atomic mass.
- (c) If the density of methanol is 0.793 kg L⁻¹, what is its volume needed for making 2.5 L of its 0.25 M solution ?
3. (a) Define molality of a solution. How is it different from molarity ?
- (b) Calculate the Molality of a solution of ethanol in water in which the mole fraction of ethanol is 0.040 (assume the density of water to be one).